

Key To

ESSENTIALS OF
TRIGONOMETRY

WELLS

QA

531

.W39

LIBRARY OF CONGRESS.

QA531
Copyright No.

Shelf..W 39

UNITED STATES OF AMERICA.

KEY


TO

WELLS' ESSENTIALS OF TRIGONOMETRY.

BY

WEBSTER WELLS, S.B.,

ASSOCIATE PROFESSOR OF MATHEMATICS IN THE MASSACHUSETTS
INSTITUTE OF TECHNOLOGY.



LEACH, SHEWELL, & SANBORN.

BOSTON AND NEW YORK.

COPYRIGHT, 1888,
BY WEBSTER WELLS.

6 A531
1039
TYPOGRAPHY BY
J. S. CUSHING & Co.

PRESSWORK BY
BERWICK & SMITH.

KEY

TO

WELLS' ESSENTIALS OF TRIGONOMETRY.

CHAPTER I.

Art. 9. — Page 3.

$$1. 135^\circ = \frac{3\pi}{4}. \qquad 5. 29^\circ 15' = \frac{13\pi}{80}.$$

$$2. 198^\circ = \frac{11\pi}{10}. \qquad 6. 174^\circ 22' 30'' = \frac{31\pi}{32}.$$

$$3. 11^\circ 15' = \frac{\pi}{16}. \qquad 7. 128^\circ 34\frac{2}{7}' = \frac{5\pi}{7}.$$

$$4. 37^\circ 30' = \frac{5\pi}{24}. \qquad 8. 92^\circ 48' 45'' = \frac{33\pi}{64}.$$

$$9. \frac{1}{2} = \frac{1}{2} \times 57.2957795^\circ \dots = 28.6478897^\circ \dots$$

$$= 28^\circ 38.873382' \dots = 28^\circ 38' 52.40292'' \dots$$

$$10. \frac{3\pi}{5} = \frac{3}{5} \text{ of } 180^\circ = 108^\circ.$$

$$11. \frac{37\pi}{30} = \frac{37}{30} \text{ of } 180^\circ = 222^\circ.$$

$$12. \frac{5\pi}{4} = \frac{5}{4} \text{ of } 180^\circ = 225^\circ.$$

$$13. \frac{3}{4} = \frac{3}{4} \times 57.2957795^\circ \dots = 42.9718346^\circ \dots$$

$$= 42^\circ 58.310076' \dots = 42^\circ 58' 18.60456'' \dots$$

$$14. 2 = 2 \times 57.2957795^\circ \dots = 114.591559^\circ \dots$$

$$= 114^\circ 35.49354' \dots = 114^\circ 35' 29.6124'' \dots$$

$$15. \quad \frac{2\pi-1}{3} = \frac{2\pi}{3} - \frac{1}{3}. \quad \frac{2\pi}{3} = \frac{2}{3} \text{ of } 180^\circ = 120^\circ.$$

$$\frac{1}{3} = \frac{1}{3} \times 57.2957795^\circ \dots = 19.0985932^\circ \dots$$

$$= 19^\circ 5.915592' \dots = 19^\circ 5' 54.93552'' \dots$$

$$\therefore \frac{2\pi-1}{3} = 120^\circ - 19^\circ 5' 54.93552'' \dots = 100^\circ 54' 5.06448'' \dots$$

$$16. \quad \frac{\pi-1}{4} = \frac{\pi}{4} - \frac{1}{4}. \quad \frac{\pi}{4} = \frac{1}{4} \text{ of } 180^\circ = 45^\circ.$$

$$\frac{1}{4} = \frac{1}{4} \times 57.2957795^\circ \dots = 14.3239449^\circ \dots$$

$$= 14^\circ 19.436694' \dots = 14^\circ 19' 26.20164'' \dots$$

$$\therefore \frac{\pi-1}{4} = 45^\circ - 14^\circ 19' 26.20164'' \dots = 30^\circ 40' 33.79836'' \dots$$

CHAPTER II.

Art. 15. — Page 8.

3. Here the opposite side = 2, and the adjacent side = 3. Therefore the hypotenuse = $\sqrt{2^2 + 3^2} = \sqrt{13}$. Then,

$$\begin{aligned}\sin A &= \frac{2}{\sqrt{13}}, & \cot A &= \frac{3}{2}, & \text{vers } A &= 1 - \frac{3}{\sqrt{13}}, \\ \cos A &= \frac{3}{\sqrt{13}}, & \sec A &= \frac{\sqrt{13}}{3}, & \text{covers } A &= 1 - \frac{2}{\sqrt{13}}, \\ & & \csc A &= \frac{\sqrt{13}}{2}, & & \end{aligned}$$

$$4. \sin A = 1 - \text{covers } A = \frac{2}{5}.$$

Here the opposite side = 2, and the hypotenuse = 5. Therefore the adjacent side = $\sqrt{5^2 - 2^2} = \sqrt{21}$. Then,

$$\begin{aligned}\cos A &= \frac{\sqrt{21}}{5}, & \cot A &= \frac{\sqrt{21}}{2}, & \csc A &= \frac{5}{2}, \\ \tan A &= \frac{2}{\sqrt{21}}, & \sec A &= \frac{5}{\sqrt{21}}, & \text{vers } A &= 1 - \frac{\sqrt{21}}{5}. \end{aligned}$$

5. Here the hypotenuse = 4, and the opposite side = 1. Therefore the adjacent side = $\sqrt{4^2 - 1^2} = \sqrt{15}$. Then,

$$\begin{aligned}\sin A &= \frac{1}{4}, & \tan A &= \frac{1}{\sqrt{15}}, & \text{vers } A &= 1 - \frac{\sqrt{15}}{4}, \\ \cos A &= \frac{\sqrt{15}}{4}, & \cot A &= \sqrt{15}, & \text{covers } A &= \frac{3}{4}, \\ & & \sec A &= \frac{4}{\sqrt{15}}, & & \end{aligned}$$

$$6. \cos A = 1 - \text{vers } A = \frac{3}{4}.$$

Here the adjacent side = 3, and the hypotenuse = 4. Therefore the opposite side = $\sqrt{4^2 - 3^2} = \sqrt{7}$. Then,

$$\begin{aligned}\sin A &= \frac{\sqrt{7}}{4}, & \cot A &= \frac{3}{\sqrt{7}}, & \csc A &= \frac{4}{\sqrt{7}}, \\ \tan A &= \frac{\sqrt{7}}{3}, & \sec A &= \frac{4}{3}, & \text{covers } A &= 1 - \frac{\sqrt{7}}{4}. \end{aligned}$$

7. Here the opposite side = x , and the hypotenuse = y . Therefore the adjacent side = $\sqrt{y^2 - x^2}$. Then,

$$\begin{aligned}\cos A &= \frac{\sqrt{y^2 - x^2}}{y}, & \cot A &= \frac{\sqrt{y^2 - x^2}}{x}, & \text{vers } A &= 1 - \frac{\sqrt{y^2 - x^2}}{y}, \\ \tan A &= \frac{x}{\sqrt{y^2 - x^2}}, & \sec A &= \frac{y}{\sqrt{y^2 - x^2}}, & \text{covers } A &= 1 - \frac{x}{y}, \\ \csc A &= \frac{y}{x},\end{aligned}$$

8. Here the hypotenuse = 13, and the adjacent side = 5. Therefore the opposite side = $\sqrt{13^2 - 5^2} = 12$. Then,

$$\begin{aligned}\sin A &= \frac{12}{13}, & \tan A &= \frac{12}{5}, & \text{vers } A &= \frac{8}{13}, \\ \cos A &= \frac{5}{13}, & \cot A &= \frac{5}{12}, & \text{covers } A &= \frac{1}{13}, \\ \csc A &= \frac{13}{12},\end{aligned}$$

9. Here the adjacent side = x , and the opposite side = 1. Therefore the hypotenuse = $\sqrt{x^2 + 1}$. Then,

$$\begin{aligned}\sin A &= \frac{1}{\sqrt{x^2 + 1}}, & \tan A &= \frac{1}{x}, & \text{vers } A &= 1 - \frac{x}{\sqrt{x^2 + 1}}, \\ \cos A &= \frac{x}{\sqrt{x^2 + 1}}, & \sec A &= \frac{\sqrt{x^2 + 1}}{x}, & \text{covers } A &= 1 - \frac{1}{\sqrt{x^2 + 1}}, \\ \csc A &= \sqrt{x^2 + 1},\end{aligned}$$

10. Here the adjacent side = 8, and the hypotenuse = 17. Therefore the opposite side = $\sqrt{17^2 - 8^2} = 15$. Then,

$$\begin{aligned}\sin A &= \frac{15}{17}, & \cot A &= \frac{8}{15}, & \text{vers } A &= \frac{9}{17}, \\ \tan A &= \frac{15}{8}, & \sec A &= \frac{17}{8}, & \text{covers } A &= \frac{2}{17}, \\ \csc A &= \frac{17}{15},\end{aligned}$$

11. Here the hypotenuse = $\sqrt{a^2 + b^2}$, and the adjacent side = b . Therefore the opposite side = $\sqrt{a^2 + b^2 - b^2} = a$. Then,

$$\begin{aligned}\sin A &= \frac{a}{\sqrt{a^2 + b^2}}, & \tan A &= \frac{a}{b}, & \text{vers } A &= 1 - \frac{b}{\sqrt{a^2 + b^2}}, \\ \cos A &= \frac{b}{\sqrt{a^2 + b^2}}, & \cot A &= \frac{b}{a}, & \text{covers } A &= 1 - \frac{a}{\sqrt{a^2 + b^2}}, \\ \csc A &= \frac{\sqrt{a^2 + b^2}}{a},\end{aligned}$$

CHAPTER III.

Art. 53. — Page 30.

3. If A is acute, $450^\circ - A$ is in the first quadrant. Then,

$$\begin{array}{ll} \sin (450^\circ - A) = \cos A, & \cos (450^\circ - A) = \sin A, \\ \tan (450^\circ - A) = \cot A, & \cot (450^\circ - A) = \tan A, \\ \sec (450^\circ - A) = \csc A, & \csc (450^\circ - A) = \sec A. \end{array}$$

4. If A is acute, $450^\circ + A$ is in the second quadrant. Then,

$$\begin{array}{ll} \sin (450^\circ + A) = \cos A, & \cos (450^\circ + A) = -\sin A, \\ \tan (450^\circ + A) = -\cot A, & \cot (450^\circ + A) = -\tan A, \\ \sec (450^\circ + A) = -\csc A, & \csc (450^\circ + A) = \sec A. \end{array}$$

5. If A is acute, $540^\circ - A$ is in the second quadrant. Then,

$$\begin{array}{ll} \sin (540^\circ - A) = \sin A, & \cos (540^\circ - A) = -\cos A, \\ \tan (540^\circ - A) = -\tan A, & \cot (540^\circ - A) = -\cot A, \\ \sec (540^\circ - A) = -\sec A, & \csc (540^\circ - A) = \csc A. \end{array}$$

6. If A is acute, $540^\circ + A$ is in the third quadrant. Then,

$$\begin{array}{ll} \sin (540^\circ + A) = -\sin A, & \cos (540^\circ + A) = -\cos A, \\ \tan (540^\circ + A) = \tan A, & \cot (540^\circ + A) = \cot A, \\ \sec (540^\circ + A) = -\sec A, & \csc (540^\circ + A) = -\csc A. \end{array}$$

7. If A is acute, $630^\circ - A$ is in the third quadrant. Then,

$$\begin{array}{ll} \sin (630^\circ - A) = -\cos A, & \cos (630^\circ - A) = -\sin A, \\ \tan (630^\circ - A) = \cot A, & \cot (630^\circ - A) = \tan A, \\ \sec (630^\circ - A) = -\csc A, & \csc (630^\circ - A) = -\sec A. \end{array}$$

8. If A is acute, $900^\circ - A$ is in the second quadrant. Then,

$$\begin{array}{ll} \sin (900^\circ - A) = \sin A, & \cos (900^\circ - A) = -\cos A, \\ \tan (900^\circ - A) = -\tan A, & \cot (900^\circ - A) = -\cot A, \\ \sec (900^\circ - A) = -\sec A, & \csc (900^\circ - A) = \csc A. \end{array}$$

9. If A is acute, $-90^\circ + A$ is in the fourth quadrant. Then,

$$\begin{array}{ll} \sin (-90^\circ + A) = -\cos A, & \cos (-90^\circ + A) = \sin A, \\ \tan (-90^\circ + A) = -\cot A, & \cot (-90^\circ + A) = -\tan A, \\ \sec (-90^\circ + A) = \csc A, & \csc (-90^\circ + A) = -\sec A. \end{array}$$

10. If A is acute, $-90^\circ - A$ is in the third quadrant. Then,
 $\sin(-90^\circ - A) = -\cos A,$ $\cos(-90^\circ - A) = -\sin A,$
 $\tan(-90^\circ - A) = \cot A,$ $\cot(-90^\circ - A) = \tan A,$
 $\sec(-90^\circ - A) = -\csc A,$ $\csc(-90^\circ - A) = -\sec A.$

11. If A is acute, $-180^\circ + A$ is in the third quadrant. Then,
 $\sin(-180^\circ + A) = -\sin A,$ $\cos(-180^\circ + A) = -\cos A,$
 $\tan(-180^\circ + A) = \tan A,$ $\cot(-180^\circ + A) = \cot A,$
 $\sec(-180^\circ + A) = -\sec A,$ $\csc(-180^\circ + A) = -\csc A.$

12. If A is acute, $-180^\circ - A$ is in the second quadrant. Then,
 $\sin(-180^\circ - A) = \sin A,$ $\cos(-180^\circ - A) = -\cos A,$
 $\tan(-180^\circ - A) = -\tan A,$ $\cot(-180^\circ - A) = -\cot A,$
 $\sec(-180^\circ - A) = -\sec A,$ $\csc(-180^\circ - A) = \csc A.$

13. If A is acute, $-270^\circ + A$ is in the second quadrant. Then,
 $\sin(-270^\circ + A) = \cos A,$ $\cos(-270^\circ + A) = -\sin A,$
 $\tan(-270^\circ + A) = -\cot A,$ $\cot(-270^\circ + A) = -\tan A,$
 $\sec(-270^\circ + A) = -\csc A,$ $\csc(-270^\circ + A) = \sec A.$

14. If A is acute, $-720^\circ + A$ is in the first quadrant. Then,
 $\sin(-720^\circ + A) = \sin A,$ $\cos(-720^\circ + A) = \cos A,$
 $\tan(-720^\circ + A) = \tan A,$ $\cot(-720^\circ + A) = \cot A,$
 $\sec(-720^\circ + A) = \sec A,$ $\csc(-720^\circ + A) = \csc A.$

Art. 54.—Page 30.

$$\begin{aligned} 2. \cos 152^\circ &= \cos(180^\circ - 28^\circ) = -\cos 28^\circ; \\ \text{or,} &= \cos(90^\circ + 62^\circ) = -\sin 62^\circ. \end{aligned}$$

$$\begin{aligned} 3. \tan 522^\circ &= \tan(540^\circ - 18^\circ) = -\tan 18^\circ; \\ \text{or,} &= \tan(450^\circ + 72^\circ) = -\cot 72^\circ. \end{aligned}$$

$$\begin{aligned} 4. \sec(-77^\circ) &= \sec(0^\circ - 77^\circ) = \sec 77^\circ; \\ \text{or,} &= \sec(-90^\circ + 13^\circ) = \csc 13^\circ. \end{aligned}$$

$$\begin{aligned} 5. \csc 230^\circ &= \csc(180^\circ + 50^\circ) = -\csc 50^\circ; \\ \text{or,} &= \csc(270^\circ - 40^\circ) = -\sec 40^\circ. \end{aligned}$$

$$\begin{aligned} 6. \cot(-129^\circ) &= \cot(-180^\circ + 51^\circ) = \cot 51^\circ; \\ \text{or,} &= \cot(-90^\circ - 39^\circ) = \tan 39^\circ. \end{aligned}$$

$$\begin{aligned}
 \text{or, } 7. \sin 865^\circ &= \sin (900^\circ - 35^\circ) = \sin 35^\circ; \\
 &= \sin (810^\circ + 55^\circ) = \cos 55^\circ. \\
 8. \cot 83^\circ &= \cot (90^\circ - 7^\circ) = \tan 7^\circ. \\
 9. \sin (-50^\circ) &= \sin (-90^\circ + 40^\circ) = -\cos 40^\circ. \\
 10. \sec 165^\circ &= \sec (180^\circ - 15^\circ) = -\sec 15^\circ. \\
 11. \cos (-303^\circ) &= \cos (-270^\circ - 33^\circ) = \sin 33^\circ. \\
 12. \tan 520^\circ &= \tan (540^\circ - 20^\circ) = -\tan 20^\circ. \\
 13. \csc 768^\circ &= \csc (810^\circ - 42^\circ) = \sec 42^\circ.
 \end{aligned}$$

Table.—Page 31.

Since $120^\circ = 180^\circ - 60^\circ$, we have

$$\begin{aligned}
 \sin 120^\circ &= \sin 60^\circ = \frac{1}{2}\sqrt{3}. & \cot 120^\circ &= -\cot 60^\circ = -\frac{1}{3}\sqrt{3}. \\
 \cos 120^\circ &= -\cos 60^\circ = -\frac{1}{2}. & \sec 120^\circ &= -\sec 60^\circ = -2. \\
 \tan 120^\circ &= -\tan 60^\circ = -\sqrt{3}. & \csc 120^\circ &= \csc 60^\circ = \frac{2}{3}\sqrt{3}.
 \end{aligned}$$

Since $135^\circ = 180^\circ - 45^\circ$, we have

$$\begin{aligned}
 \sin 135^\circ &= \sin 45^\circ = \frac{1}{2}\sqrt{2}. & \cot 135^\circ &= -\cot 45^\circ = -1. \\
 \cos 135^\circ &= -\cos 45^\circ = -\frac{1}{2}\sqrt{2}. & \sec 135^\circ &= -\sec 45^\circ = -\sqrt{2}. \\
 \tan 135^\circ &= -\tan 45^\circ = -1. & \csc 135^\circ &= \csc 45^\circ = \sqrt{2}.
 \end{aligned}$$

Since $150^\circ = 180^\circ - 30^\circ$, we have

$$\begin{aligned}
 \sin 150^\circ &= \sin 30^\circ = \frac{1}{2}. & \cot 150^\circ &= -\cot 30^\circ = -\sqrt{3}. \\
 \cos 150^\circ &= -\cos 30^\circ = -\frac{1}{2}\sqrt{3}. & \sec 150^\circ &= -\sec 30^\circ = -\frac{2}{3}\sqrt{3}. \\
 \tan 150^\circ &= -\tan 30^\circ = -\frac{1}{3}\sqrt{3}. & \csc 150^\circ &= \csc 30^\circ = 2.
 \end{aligned}$$

Since $210^\circ = 180^\circ + 30^\circ$, we have

$$\begin{aligned}
 \sin 210^\circ &= -\sin 30^\circ = -\frac{1}{2}. & \cot 210^\circ &= \cot 30^\circ = \sqrt{3}. \\
 \cos 210^\circ &= -\cos 30^\circ = -\frac{1}{2}\sqrt{3}. & \sec 210^\circ &= -\sec 30^\circ = -\frac{2}{3}\sqrt{3}. \\
 \tan 210^\circ &= \tan 30^\circ = \frac{1}{3}\sqrt{3}. & \csc 210^\circ &= -\csc 30^\circ = -2.
 \end{aligned}$$

Since $225^\circ = 180^\circ + 45^\circ$, we have

$$\begin{aligned}\sin 225^\circ &= -\sin 45^\circ = -\frac{1}{2}\sqrt{2}. & \cot 225^\circ &= \cot 45^\circ = 1. \\ \cos 225^\circ &= -\cos 45^\circ = -\frac{1}{2}\sqrt{2}. & \sec 225^\circ &= -\sec 45^\circ = -\sqrt{2}. \\ \tan 225^\circ &= \tan 45^\circ = 1. & \csc 225^\circ &= -\csc 45^\circ = -\sqrt{2}.\end{aligned}$$

Since $240^\circ = 180^\circ + 60^\circ$, we have

$$\begin{aligned}\sin 240^\circ &= -\sin 60^\circ = -\frac{1}{2}\sqrt{3}. & \cot 240^\circ &= \cot 60^\circ = \frac{1}{3}\sqrt{3}. \\ \cos 240^\circ &= -\cos 60^\circ = -\frac{1}{2}. & \sec 240^\circ &= -\sec 60^\circ = -2. \\ \tan 240^\circ &= \tan 60^\circ = \sqrt{3}. & \csc 240^\circ &= -\csc 60^\circ = -\frac{2}{3}\sqrt{3}.\end{aligned}$$

Since $300^\circ = 360^\circ - 60^\circ$, we have

$$\begin{aligned}\sin 300^\circ &= -\sin 60^\circ = -\frac{1}{2}\sqrt{3}. & \cot 300^\circ &= -\cot 60^\circ = -\frac{1}{3}\sqrt{3}. \\ \cos 300^\circ &= \cos 60^\circ = \frac{1}{2}. & \sec 300^\circ &= \sec 60^\circ = 2. \\ \tan 300^\circ &= -\tan 60^\circ = -\sqrt{3}. & \csc 300^\circ &= -\csc 60^\circ = -\frac{2}{3}\sqrt{3}.\end{aligned}$$

Since $315^\circ = 360^\circ - 45^\circ$, we have

$$\begin{aligned}\sin 315^\circ &= -\sin 45^\circ = -\frac{1}{2}\sqrt{2}. & \cot 315^\circ &= -\cot 45^\circ = -1. \\ \cos 315^\circ &= \cos 45^\circ = \frac{1}{2}\sqrt{2}. & \sec 315^\circ &= \sec 45^\circ = \sqrt{2}. \\ \tan 315^\circ &= -\tan 45^\circ = -1. & \csc 315^\circ &= -\csc 45^\circ = -\sqrt{2}.\end{aligned}$$

Since $330^\circ = 360^\circ - 30^\circ$, we have

$$\begin{aligned}\sin 330^\circ &= -\sin 30^\circ = -\frac{1}{2}. & \cot 330^\circ &= -\cot 30^\circ = -\sqrt{3}. \\ \cos 330^\circ &= \cos 30^\circ = \frac{1}{2}\sqrt{3}. & \sec 330^\circ &= \sec 30^\circ = \frac{2}{3}\sqrt{3}. \\ \tan 330^\circ &= -\tan 30^\circ = -\frac{1}{3}\sqrt{3}. & \csc 330^\circ &= -\csc 30^\circ = -2.\end{aligned}$$

Art. 56.—Page 34.

3. Here the ordinate = 1, and the distance = 4. Therefore the abscissa = $\pm\sqrt{4^2-1^2} = \pm\sqrt{15}$. Then,

$$\cos A = \pm \frac{\sqrt{15}}{4}, \quad \cot A = \pm \sqrt{15}, \quad \csc A = 4.$$

$$\tan A = \pm \frac{1}{\sqrt{15}}, \quad \sec A = \pm \frac{4}{\sqrt{15}},$$

4. Here the abscissa = 2 and the ordinate = 1, or the abscissa = -2 and the ordinate = -1. Therefore the distance = $\sqrt{2^2+1^2} = \sqrt{5}$. Then,

$$\sin A = \pm \frac{1}{\sqrt{5}}, \quad \tan A = \frac{1}{2}, \quad \csc A = \pm \sqrt{5}.$$

$$\cos A = \pm \frac{2}{\sqrt{5}}, \quad \sec A = \pm \frac{\sqrt{5}}{2},$$

5. Here the distance = 3, and the ordinate = -2. Therefore the abscissa = $\pm\sqrt{3^2-2^2} = \pm\sqrt{5}$. Then,

$$\sin A = -\frac{2}{3}, \quad \tan A = \mp \frac{2}{\sqrt{5}}, \quad \sec A = \pm \frac{3}{\sqrt{5}}.$$

$$\cos A = \pm \frac{\sqrt{5}}{3}, \quad \cot A = \mp \frac{\sqrt{5}}{2},$$

6. Here the ordinate = 8 and the abscissa = -15, or the ordinate = -8 and the abscissa = 15. Therefore the distance = $\sqrt{8^2+15^2} = 17$. Then,

$$\sin A = \pm \frac{8}{17}, \quad \cot A = -\frac{15}{8}, \quad \csc A = \pm \frac{17}{8}.$$

$$\cos A = \mp \frac{15}{17}, \quad \sec A = \mp \frac{17}{15},$$

7. Here the distance = 4, and the abscissa = 3. Therefore the ordinate = $\pm\sqrt{4^2-3^2} = \pm\sqrt{7}$. Then,

$$\sin A = \pm \frac{\sqrt{7}}{4}, \quad \tan A = \pm \frac{\sqrt{7}}{3}, \quad \csc A = \pm \frac{4}{\sqrt{7}}.$$

$$\cos A = \frac{3}{4}, \quad \cot A = \pm \frac{3}{\sqrt{7}},$$

8. Here the abscissa = -1, and the distance = 2. Therefore the ordinate = $\pm\sqrt{2^2-1^2} = \pm\sqrt{3}$. Then,

$$\sin A = \pm \frac{\sqrt{3}}{2}, \quad \cot A = \mp \frac{1}{\sqrt{3}}, \quad \csc A = \pm \frac{2}{\sqrt{3}}.$$

$$\tan A = \mp \sqrt{3}, \quad \sec A = -2,$$

9. Here the distance = $\sqrt{2}$, and the ordinate = 1. Therefore the abscissa = $\pm \sqrt{2-1} = \pm 1$. Then,

$$\sin A = \frac{1}{\sqrt{2}}, \quad \tan A = \pm 1, \quad \sec A = \pm \sqrt{2}.$$

$$\cos A = \pm \frac{1}{\sqrt{2}}, \quad \cot A = \pm 1,$$

10. Here the ordinate = $2\sqrt{2}$ and the abscissa = 1, or the ordinate = $-2\sqrt{2}$ and the abscissa = -1. Therefore the distance = $\sqrt{8+1} = 3$. Then,

$$\sin A = \pm \frac{2\sqrt{2}}{3}, \quad \cot A = \frac{1}{2\sqrt{2}}, \quad \csc A = \pm \frac{3}{2\sqrt{2}}.$$

$$\cos A = \pm \frac{1}{3}, \quad \sec A = \pm 3,$$

11. Here the abscissa = $-a$, and the distance = b . Therefore the ordinate = $\pm \sqrt{b^2 - a^2}$. Then,

$$\sin A = \pm \frac{\sqrt{b^2 - a^2}}{b}, \quad \cot A = \mp \frac{a}{\sqrt{b^2 - a^2}}, \quad \csc A = \pm \frac{b}{\sqrt{b^2 - a^2}}.$$

$$\tan A = \mp \frac{\sqrt{b^2 - a^2}}{a}, \quad \sec A = -\frac{b}{a},$$

12. Here the ordinate = x , and the distance = 1. Therefore the abscissa = $\pm \sqrt{1 - x^2}$. Then,

$$\cos A = \pm \sqrt{1 - x^2}, \quad \cot A = \pm \frac{\sqrt{1 - x^2}}{x}, \quad \csc A = \frac{1}{x}.$$

$$\tan A = \pm \frac{x}{\sqrt{1 - x^2}}, \quad \sec A = \pm \frac{1}{\sqrt{1 - x^2}},$$

13. Here the abscissa = 1 and the ordinate = x , or the abscissa = -1 and the ordinate = $-x$. Therefore the distance = $\sqrt{1 + x^2}$. Then,

$$\sin A = \pm \frac{x}{\sqrt{1 + x^2}}, \quad \tan A = x, \quad \csc A = \pm \frac{\sqrt{1 + x^2}}{x}.$$

$$\cos A = \pm \frac{1}{\sqrt{1 + x^2}}, \quad \sec A = \pm \sqrt{1 + x^2},$$

14. Here the distance = $\sqrt{a^2 + b^2}$, and the abscissa = a . Therefore the ordinate = $\pm \sqrt{a^2 + b^2 - a^2} = \pm b$. Then,

$$\sin A = \pm \frac{b}{\sqrt{a^2 + b^2}}, \quad \tan A = \pm \frac{b}{a}, \quad \csc A = \pm \frac{\sqrt{a^2 + b^2}}{b}.$$

$$\cos A = \frac{a}{\sqrt{a^2 + b^2}}, \quad \cot A = \pm \frac{a}{b},$$

CHAPTER IV.

Art. 60. — Page 36.

$$\sin A = \tan A \cos A \text{ (Art. 59)} = \frac{\tan A}{\sec A} = \frac{\tan A}{\sqrt{1 + \tan^2 A}} \text{ (Art. 58).}$$

$$\sin A = \frac{1}{\csc A} = \frac{1}{\sqrt{1 + \cot^2 A}} \text{ (Art. 58).}$$

$$\sin A = \sqrt{1 - \cos^2 A} = \sqrt{1 - \frac{1}{\sec^2 A}} = \frac{\sqrt{\sec^2 A - 1}}{\sec A}.$$

$$\cos A = \frac{1}{\sec A} = \frac{1}{\sqrt{1 + \tan^2 A}} \text{ (Art. 58).}$$

$$\cos A = \cot A \sin A \text{ (Art. 59)} = \frac{\cot A}{\csc A} = \frac{\cot A}{\sqrt{1 + \cot^2 A}} \text{ (Art. 58).}$$

$$\cos A = \sqrt{1 - \sin^2 A} = \sqrt{1 - \frac{1}{\csc^2 A}} = \frac{\sqrt{\csc^2 A - 1}}{\csc A}.$$

$$\tan A = \frac{\sin A}{\cos A} = \frac{\sin A}{\sqrt{1 - \sin^2 A}} = \frac{\sqrt{1 - \cos^2 A}}{\cos A}.$$

$$\tan A = \frac{1}{\cot A} = \frac{1}{\sqrt{\csc^2 A - 1}} \text{ (Art. 58).}$$

Since the cotangent, secant, and cosecant are the reciprocals of the tangent, cosine, and sine, respectively, we have :

$$\cot A = \frac{\sqrt{1 - \sin^2 A}}{\sin A} = \frac{\cos A}{\sqrt{1 - \cos^2 A}} = \frac{1}{\sqrt{\sec^2 A - 1}}.$$

$$\sec A = \frac{1}{\sqrt{1 - \sin^2 A}} = \frac{\sqrt{1 + \cot^2 A}}{\cot A} = \frac{\csc A}{\sqrt{\csc^2 A - 1}}.$$

$$\csc A = \frac{1}{\sqrt{1 - \cos^2 A}} = \frac{\sqrt{1 + \tan^2 A}}{\tan A} = \frac{\sec A}{\sqrt{\sec^2 A - 1}}.$$

CHAPTER V.

Art. 78. — Pages 51 to 53.

- $$\begin{aligned}
 3. \quad \frac{\sin(x+y)}{\sin(x-y)} &= \frac{\sin x \cos y + \cos x \sin y}{\sin x \cos y - \cos x \sin y} \\
 &= \frac{\frac{\sin x \cos y}{\cos x \cos y} + \frac{\cos x \sin y}{\cos x \cos y}}{\frac{\sin x \cos y}{\cos x \cos y} - \frac{\cos x \sin y}{\cos x \cos y}} = \frac{\tan x + \tan y}{\tan x - \tan y}.
 \end{aligned}$$
- $$\begin{aligned}
 4. \quad \frac{\cos(x+y)}{\cos(x-y)} &= \frac{\cos x \cos y - \sin x \sin y}{\cos x \cos y + \sin x \sin y} \\
 &= \frac{\frac{\cos x \cos y}{\sin x \sin y} - \frac{\sin x \sin y}{\sin x \sin y}}{\frac{\cos x \cos y}{\sin x \sin y} + \frac{\sin x \sin y}{\sin x \sin y}} = \frac{\cot x \cot y - 1}{\cot x \cot y + 1}.
 \end{aligned}$$
- $$\begin{aligned}
 5. \quad \frac{\sin(x+y)}{\cos(x-y)} &= \frac{\sin x \cos y + \cos x \sin y}{\cos x \cos y + \sin x \sin y} \\
 &= \frac{\frac{\sin x \cos y}{\sin x \cos y} + \frac{\cos x \sin y}{\sin x \cos y}}{\frac{\cos x \cos y}{\sin x \cos y} + \frac{\sin x \sin y}{\sin x \cos y}} = \frac{1 + \cot x \tan y}{\cot x + \tan y}.
 \end{aligned}$$
- $$\begin{aligned}
 6. \quad \sin(45^\circ + y) &= \sin 45^\circ \cos y + \cos 45^\circ \sin y \\
 &= \frac{1}{\sqrt{2}} \cdot \cos y + \frac{1}{\sqrt{2}} \cdot \sin y \quad (\text{Art. 16}) = \frac{\sin y + \cos y}{\sqrt{2}}.
 \end{aligned}$$
- $$7. \quad \tan(60^\circ - y) = \frac{\tan 60^\circ - \tan y}{1 + \tan 60^\circ \tan y} = \frac{\sqrt{3} - \tan y}{1 + \sqrt{3} \tan y} \quad (\text{Art. 17}).$$
- $$8. \quad \frac{\sin x + \sin y}{\cos x + \cos y} = \frac{2 \sin \frac{1}{2}(x+y) \cos \frac{1}{2}(x-y)}{2 \cos \frac{1}{2}(x+y) \cos \frac{1}{2}(x-y)} = \tan \frac{1}{2}(x+y).$$
- $$9. \quad \frac{\sin x + \sin y}{\cos x - \cos y} = \frac{2 \sin \frac{1}{2}(x+y) \cos \frac{1}{2}(x-y)}{-2 \sin \frac{1}{2}(x+y) \sin \frac{1}{2}(x-y)} = -\cot \frac{1}{2}(x-y).$$

10. By Arts. 74 and 58, $\sin 2x = \frac{2 \sin x \cos x}{\sin^2 x + \cos^2 x}$.

Dividing each term of the fraction by $\cos^2 x$,

$$\sin 2x = \frac{\frac{2 \sin x}{\cos x}}{1 + \frac{\sin^2 x}{\cos^2 x}} = \frac{2 \tan x}{1 + \tan^2 x}.$$

11. $\csc 2x = \frac{1}{\sin 2x} = \frac{1}{2 \sin x \cos x} = \frac{1}{2} \sec x \csc x$ (Art. 57).

12. $\tan x + \cot x = \frac{\sin x}{\cos x} + \frac{\cos x}{\sin x} = \frac{\sin^2 x + \cos^2 x}{\sin x \cos x}$
 $= \frac{1}{\sin x \cos x} = \frac{2}{2 \sin x \cos x} = \frac{2}{\sin 2x}.$

13. $\cot x - \tan x = \cot x - \frac{1}{\cot x} = \frac{\cot^2 x - 1}{\cot x}$
 $= 2 \left(\frac{\cot^2 x - 1}{2 \cot x} \right) = 2 \cot 2x$ (Art. 74).

14. $\frac{(1 + \tan x)^2 - (1 - \tan x)^2}{(1 + \tan x)^2 + (1 - \tan x)^2}$
 $= \frac{1 + 2 \tan x + \tan^2 x - 1 + 2 \tan x - \tan^2 x}{1 + 2 \tan x + \tan^2 x + 1 - 2 \tan x + \tan^2 x}$
 $= \frac{4 \tan x}{2 + 2 \tan^2 x} = \frac{2 \tan x}{1 + \tan^2 x} = \sin 2x$, by Ex. 10

15. $\sin(x + y) \sin(x - y)$
 $= (\sin x \cos y + \cos x \sin y)(\sin x \cos y - \cos x \sin y)$
 $= \sin^2 x \cos^2 y - \cos^2 x \sin^2 y$
 $= \sin^2 x (1 - \sin^2 y) - (1 - \sin^2 x) \sin^2 y$
 $= \sin^2 x - \sin^2 y.$

16. $\cos(x + y) \cos(x - y)$
 $= (\cos x \cos y - \sin x \sin y)(\cos x \cos y + \sin x \sin y)$
 $= \cos^2 x \cos^2 y - \sin^2 x \sin^2 y$
 $= \cos^2 x \cos^2 y - (1 - \cos^2 x)(1 - \cos^2 y)$
 $= \cos^2 x \cos^2 y - 1 + \cos^2 x + \cos^2 y - \cos^2 x \cos^2 y$
 $= \cos^2 x - (1 - \cos^2 y) = \cos^2 x - \sin^2 y.$

$$\begin{aligned}
 17. \sec^2 x \csc^2 x &= \frac{1}{\cos^2 x \sin^2 x} = \frac{\sin^2 x + \cos^2 x}{\cos^2 x \sin^2 x} \\
 &= \frac{\sin^2 x}{\cos^2 x \sin^2 x} + \frac{\cos^2 x}{\cos^2 x \sin^2 x} = \frac{1}{\cos^2 x} + \frac{1}{\sin^2 x} \\
 &= \sec^2 x + \csc^2 x.
 \end{aligned}$$

$$\begin{aligned}
 18. \cos y + \cos (120^\circ + y) + \cos (120^\circ - y) \\
 = \cos y + \cos 120^\circ \cos y - \sin 120^\circ \sin y + \cos 120^\circ \cos y + \sin 120^\circ \sin y \\
 = \cos y + 2 \cos 120^\circ \cos y = \cos y - \cos y \text{ (Art. 55)} = 0.
 \end{aligned}$$

$$\begin{aligned}
 19. \sin A \sin (B - C) + \sin B \sin (C - A) + \sin C \sin (A - B) \\
 = \sin A (\sin B \cos C - \cos B \sin C) + \sin B (\sin C \cos A - \cos C \sin A) \\
 + \sin C (\sin A \cos B - \cos A \sin B) \\
 = \sin A \sin B \cos C - \sin A \cos B \sin C + \sin B \sin C \cos A \\
 - \sin B \cos C \sin A + \sin C \sin A \cos B - \sin C \cos A \sin B = 0.
 \end{aligned}$$

$$\begin{aligned}
 20. \cos (A + B) \cos (A - B) + \cos (B + C) \cos (B - C) \\
 + \cos (C + A) \cos (C - A) \\
 = \cos^2 A - \sin^2 B + \cos^2 B - \sin^2 C + \cos^2 C - \sin^2 A, \text{ by Ex. 16,} \\
 = \cos 2A + \cos 2B + \cos 2C \text{ (Art. 74).}
 \end{aligned}$$

$$\begin{aligned}
 21. \frac{\cos x - \cos 3x}{\sin 3x - \sin x} &= \frac{-(\cos 3x - \cos x)}{\sin 3x - \sin x} \\
 &= \frac{2 \sin \frac{1}{2}(3x + x) \sin \frac{1}{2}(3x - x)}{2 \cos \frac{1}{2}(3x + x) \sin \frac{1}{2}(3x - x)} = \tan 2x.
 \end{aligned}$$

$$\begin{aligned}
 22. \frac{\cos 80^\circ + \cos 20^\circ}{\sin 80^\circ - \sin 20^\circ} &= \frac{2 \cos \frac{1}{2}(80^\circ + 20^\circ) \cos \frac{1}{2}(80^\circ - 20^\circ)}{2 \cos \frac{1}{2}(80^\circ + 20^\circ) \sin \frac{1}{2}(80^\circ - 20^\circ)} \\
 &= \cot 30^\circ = \sqrt{3} \text{ (Art. 17).}
 \end{aligned}$$

$$\begin{aligned}
 23. \sin (x + y + z) &= \sin [(x + y) + z] \\
 &= \sin (x + y) \cos z + \cos (x + y) \sin z \\
 &= (\sin x \cos y + \cos x \sin y) \cos z + (\cos x \cos y - \sin x \sin y) \sin z \\
 &= \sin x \cos y \cos z + \cos x \sin y \cos z + \cos x \cos y \sin z - \sin x \sin y \sin z.
 \end{aligned}$$

$$\begin{aligned}
 24. \cos (x + y + z) &= \cos [(x + y) + z] \\
 &= \cos (x + y) \cos z - \sin (x + y) \sin z \\
 &= (\cos x \cos y - \sin x \sin y) \cos z - (\sin x \cos y + \cos x \sin y) \sin z \\
 &= \cos x \cos y \cos z - \sin x \sin y \cos z - \sin x \cos y \sin z - \cos x \sin y \sin z.
 \end{aligned}$$

$$\begin{aligned}
 25. \quad \sin 3x &= \sin (2x + x) = \sin 2x \cos x + \cos 2x \sin x \\
 &= 2 \sin x \cos^2 x + (1 - 2 \sin^2 x) \sin x \quad (\text{Art. 74}) \\
 &= 2 \sin x (1 - \sin^2 x) + \sin x - 2 \sin^3 x \\
 &= 3 \sin x - 4 \sin^3 x.
 \end{aligned}$$

$$\begin{aligned}
 26. \quad \cos 3x &= \cos (2x + x) = \cos 2x \cos x - \sin 2x \sin x \\
 &= (2 \cos^2 x - 1) \cos x - 2 \sin^2 x \cos x \quad (\text{Art. 74}) \\
 &= 2 \cos^3 x - \cos x - 2 (1 - \cos^2 x) \cos x \\
 &= 4 \cos^3 x - 3 \cos x.
 \end{aligned}$$

$$\begin{aligned}
 27. \quad \tan 3x &= \tan (2x + x) = \frac{\tan 2x + \tan x}{1 - \tan 2x \tan x} \\
 &= \frac{\frac{2 \tan x}{1 - \tan^2 x} + \tan x}{1 - \frac{2 \tan^2 x}{1 - \tan^2 x}} \quad (\text{Art. 74}) \\
 &= \frac{2 \tan x + \tan x - \tan^3 x}{1 - \tan^2 x - 2 \tan^2 x} = \frac{3 \tan x - \tan^3 x}{1 - 3 \tan^2 x}.
 \end{aligned}$$

$$\begin{aligned}
 28. \quad \sin (2x + y) - 2 \sin x \cos (x + y) \\
 &= \sin [(x + y) + x] - 2 \sin x \cos (x + y) \\
 &= \sin (x + y) \cos x + \cos (x + y) \sin x - 2 \sin x \cos (x + y) \\
 &= \sin (x + y) \cos x - \cos (x + y) \sin x \\
 &= \sin [(x + y) - x] = \sin y.
 \end{aligned}$$

$$\begin{aligned}
 29. \quad \frac{\sin 3x}{\sin x} - \frac{\cos 3x}{\cos x} &= \frac{\sin 3x \cos x - \cos 3x \sin x}{\sin x \cos x} \\
 &= \frac{\sin (3x - x)}{\frac{1}{2} \sin 2x} \quad (\text{Art. 74}) = 2.
 \end{aligned}$$

$$\begin{aligned}
 30. \quad 1 + \cos 2x \cos 2y &= 1 + (2 \cos^2 x - 1)(1 - 2 \sin^2 y) \\
 &= 2 \cos^2 x + 2 \sin^2 y - 4 \cos^2 x \sin^2 y \\
 &= 2 \cos^2 x (\sin^2 y + \cos^2 y) + 2 \sin^2 y (\sin^2 x + \cos^2 x) - 4 \cos^2 x \sin^2 y \\
 &= 2 (\sin^2 x \sin^2 y + \cos^2 x \cos^2 y).
 \end{aligned}$$

$$\begin{aligned}
 31. \quad 1 + \tan x \tan 2x &= 1 + \frac{\sin x \sin 2x}{\cos x \cos 2x} \\
 &= \frac{\cos 2x \cos x + \sin 2x \sin x}{\cos 2x \cos x} = \frac{\cos (2x - x)}{\cos 2x \cos x} = \frac{1}{\cos 2x} = \sec 2x.
 \end{aligned}$$

$$32. \sin 4x = 2 \sin 2x \cos 2x = 4 \sin x \cos x (1 - 2 \sin^2 x) \text{ (Art. 74)} \\ = 4 \sin x \cos x - 8 \sin^3 x \cos x.$$

$$33. \cos 4x = 2 \cos^2 2x - 1 = 2(2 \cos^2 x - 1)^2 - 1 \text{ (Art. 74)} \\ = 8 \cos^4 x - 8 \cos^2 x + 2 - 1 = 8 \cos^4 x - 8 \cos^2 x + 1.$$

$$34. \sin 5x = \sin (4x + x) = \sin 4x \cos x + \cos 4x \sin x \\ = 4 \sin x \cos^2 x - 8 \sin^3 x \cos^2 x + 8 \cos^4 x \sin x - 8 \cos^2 x \sin x \\ + \sin x \text{ (Exs. 32, 33)} \\ = 4 \sin x (1 - \sin^2 x) - 8 \sin^3 x (1 - \sin^2 x) + 8 (1 - \sin^2 x)^2 \sin x \\ - 8 (1 - \sin^2 x) \sin x + \sin x \\ = 4 \sin x - 4 \sin^3 x - 8 \sin^3 x + 8 \sin^5 x + 8 \sin x - 16 \sin^3 x \\ + 8 \sin^5 x - 8 \sin x + 8 \sin^3 x + \sin x \\ = 5 \sin x - 20 \sin^3 x + 16 \sin^5 x.$$

$$35. \sin 15^\circ = \sin (45^\circ - 30^\circ) = \sin 45^\circ \cos 30^\circ - \cos 45^\circ \sin 30^\circ \\ = \frac{1}{2} \sqrt{2} \cdot \frac{1}{2} \sqrt{3} - \frac{1}{2} \sqrt{2} \cdot \frac{1}{2} = \frac{1}{4} (\sqrt{6} - \sqrt{2}) = \cos 75^\circ. \\ \cos 15^\circ = \cos (45^\circ - 30^\circ) = \cos 45^\circ \cos 30^\circ + \sin 45^\circ \sin 30^\circ \\ = \frac{1}{2} \sqrt{2} \cdot \frac{1}{2} \sqrt{3} + \frac{1}{2} \sqrt{2} \cdot \frac{1}{2} = \frac{1}{4} (\sqrt{6} + \sqrt{2}) = \sin 75^\circ.$$

$$36. \tan 15^\circ = \frac{1 - \cos 30^\circ}{\sin 30^\circ} = \csc 30^\circ - \cot 30^\circ = 2 - \sqrt{3} = \cot 75^\circ. \\ \cot 15^\circ = \frac{1 + \cos 30^\circ}{\sin 30^\circ} = \csc 30^\circ + \cot 30^\circ = 2 + \sqrt{3} = \tan 75^\circ.$$

$$37. \sin 22^\circ 30' = \sqrt{\frac{1 - \cos 45^\circ}{2}} = \sqrt{\frac{1 - \frac{1}{2} \sqrt{2}}{2}} = \sqrt{\frac{2 - \sqrt{2}}{4}} \\ = \frac{1}{2} \sqrt{2 - \sqrt{2}}. \\ \cos 22^\circ 30' = \sqrt{\frac{1 + \cos 45^\circ}{2}} = \sqrt{\frac{1 + \frac{1}{2} \sqrt{2}}{2}} = \sqrt{\frac{2 + \sqrt{2}}{4}} \\ = \frac{1}{2} \sqrt{2 + \sqrt{2}}.$$

$$38. \tan 22^\circ 30' = \frac{1 - \cos 45^\circ}{\sin 45^\circ} = \csc 45^\circ - \cot 45^\circ = \sqrt{2} - 1. \\ \cot 22^\circ 30' = \frac{1 + \cos 45^\circ}{\sin 45^\circ} = \csc 45^\circ + \cot 45^\circ = \sqrt{2} + 1.$$

CHAPTER VI.

Art. 91.—Page 57.

2. $\log 6 = \log (2 \times 3) = \log 2 + \log 3 = .3010 + .4771 = .7781.$
3. $\log 14 = \log (2 \times 7) = \log 2 + \log 7 = .3010 + .8451 = 1.1461.$
4. $\log 8 = \log (2 \times 2 \times 2) = \log 2 + \log 2 + \log 2 = 3 \log 2$
 $= 3 \times .3010 = .9030.$
5. $\log 12 = \log (2 \times 2 \times 3) = \log 2 + \log 2 + \log 3$
 $= 2 \log 2 + \log 3 = .6020 + .4771 = 1.0791.$
6. $\log 15 = \log (3 \times 5) = \log 3 + \log 5 = .4771 + .6990 = 1.1761.$
7. $\log 21 = \log (3 \times 7) = \log 3 + \log 7 = .4771 + .8451 = 1.3222.$
8. $\log 63 = \log (3 \times 3 \times 7) = \log 3 + \log 3 + \log 7 = 2 \log 3 + \log 7$
 $= .9542 + .8451 = 1.7993.$
9. $\log 56 = \log (2 \times 2 \times 2 \times 7) = 3 \log 2 + \log 7$
 $= .9030 + .8451 = 1.7481.$
10. $\log 84 = \log (2 \times 2 \times 3 \times 7) = 2 \log 2 + \log 3 + \log 7$
 $= .6020 + .4771 + .8451 = 1.9242.$
11. $\log 45 = \log (3 \times 3 \times 5) = 2 \log 3 + \log 5 = .9542 + .6990 = 1.6532.$
12. $\log 98 = \log (2 \times 7 \times 7) = \log 2 + 2 \log 7 = .3010 + 1.6902 = 1.9912.$
13. $\log 105 = \log (3 \times 5 \times 7) = \log 3 + \log 5 + \log 7$
 $= .4771 + .6990 + .8451 = 2.0212.$
14. $\log 112 = \log (2 \times 2 \times 2 \times 2 \times 7) = 4 \log 2 + \log 7$
 $= 1.2040 + .8451 = 2.0491.$
15. $\log 144 = \log (2 \times 2 \times 2 \times 2 \times 3 \times 3) = 4 \log 2 + 2 \log 3$
 $= 1.2040 + .9542 = 2.1582.$
16. $\log 216 = \log (2 \times 2 \times 2 \times 3 \times 3 \times 3) = 3 \log 2 + 3 \log 3$
 $= .9030 + 1.4313 = 2.3343.$

17. $\log 135 = \log (3 \times 3 \times 3 \times 5) = 3 \log 3 + \log 5$
 $= 1.4313 + .6990 = 2.1303.$
18. $\log 168 = \log (2 \times 2 \times 2 \times 3 \times 7) = 3 \log 2 + \log 3 + \log 7$
 $= .9030 + .4771 + .8451 = 2.2252.$
19. $\log 147 = \log (3 \times 7 \times 7) = \log 3 + 2 \log 7 = .4771 + 1.6902 = 2.1673.$
20. $\log 375 = \log (3 \times 5 \times 5 \times 5) = \log 3 + 3 \log 5$
 $= .4771 + 2.0970 = 2.5741.$
21. $\log 343 = \log (7 \times 7 \times 7) = 3 \log 7 = 2.5353.$

Art. 93.—Page 58.

2. $\log \frac{7}{3} = \log 7 - \log 3 = .8451 - .4771 = .3680.$
3. $\log \frac{10}{7} = \log 10 - \log 7 = 1 - .8451 = .1549.$
4. $\log 3\frac{1}{3} = \log \frac{10}{3} = \log 10 - \log 3 = 1 - .4771 = .5229.$
5. $\log 35 = \log \frac{70}{2} = \log (10 \times 7) - \log 2 = \log 10 + \log 7 - \log 2$
 $= 1 + .8451 - .3010 = 1.5441.$
6. $\log \frac{21}{16} = \log 21 - \log 16 = \log (3 \times 7) - \log (2 \times 2 \times 2 \times 2)$
 $= \log 3 + \log 7 - 4 \log 2 = .4771 + .8451 - 1.2040 = .1182.$
7. $\log 125 = \log (5 \times 5 \times 5) = 3 \log 5.$
 $\log 5 = \log \frac{10}{2} = 1 - \log 2 = 1 - .3010 = .6990.$
 $\therefore \log 125 = 3 \times .6990 = 2.0970.$
8. $\log \frac{42}{25} = \log (2 \times 3 \times 7) - \log (5 \times 5) = \log 2 + \log 3 + \log 7 - 2 \log 5.$
 $\log 5 = \log \frac{10}{2} = 1 - \log 2 = .6990.$
 $\therefore \log \frac{42}{25} = .3010 + .4771 + .8451 - 1.3980 = .2252.$
9. $\log 175 = \log (5 \times 5 \times 7) = 2 \log 5 + \log 7.$
 $\log 5 = \log \frac{10}{2} = 1 - \log 2 = .6990.$
 $\therefore \log 175 = 1.3980 + .8451 = 2.2431.$

10. $\log 11\frac{1}{3} = \log \frac{100}{9} = \log 100 - \log (3 \times 3)$
 $= 2 - 2 \log 3 = 2 - .9542 = 1.0458.$
11. $\log 7\frac{1}{7} = \log \frac{50}{7} = \log \frac{100}{14} = \log 100 - \log (2 \times 7)$
 $= 2 - \log 2 - \log 7 = 2 - .3010 - .8451 = .8539.$
12. $\log \frac{35}{6} = \log (5 \times 7) - \log (2 \times 3) = \log 5 + \log 7 - \log 2 - \log 3.$
 $\log 5 = \log \frac{10}{2} = 1 - \log 2 = .6990.$
 $\therefore \log \frac{35}{6} = .6990 + .8451 - .3010 - .4771 = .7660.$
13. $\log 5\frac{4}{9} = \log \frac{49}{9} = \log (7 \times 7) - \log (3 \times 3) = 2 \log 7 - 2 \log 3$
 $= 1.6902 - .9542 = .7360.$

Art. 96.—Page 59.

3. $\log 3^{\frac{3}{5}} = \frac{3}{5} \log 3 = \frac{3}{5} \times .4771 = .2863.$
4. $\log 2^9 = 9 \log 2 = 9 \times .3010 = 2.7090.$
5. $\log 7^5 = 5 \log 7 = 5 \times .8451 = 4.2255.$
6. $\log 5^{\frac{1}{5}} = \frac{1}{5} \log 5.$
 $\log 5 = \log \frac{10}{2} = 1 - \log 2 = .6990.$
 $\therefore \log 5^{\frac{1}{5}} = \frac{.6990}{5} = .1398.$
7. $\log 12^{\frac{2}{3}} = \frac{2}{3} \log 12.$
 $\log 12 = \log (2 \times 2 \times 3) = 2 \log 2 + \log 3 = .6020 + .4771 = 1.0791.$
 $\therefore \log 12^{\frac{2}{3}} = \frac{2}{3} \times 1.0791 = .7194.$
8. $\log 21^{\frac{1}{2}} = \frac{1}{2} \log 21.$
 $\log 21 = \log (3 \times 7) = \log 3 + \log 7 = .4771 + .8451 = 1.3222.$
 $\therefore \log 21^{\frac{1}{2}} = \frac{1.3222}{2} = .6611.$

$$9. \log 14^4 = 4 \log 14.$$

$$\log 14 = \log (2 \times 7) = \log 2 + \log 7 = .3010 + .8451 = 1.1461.$$

$$\therefore \log 14^4 = 4 \times 1.1461 = 4.5844.$$

$$10. \log 25^{\frac{7}{3}} = \frac{7}{3} \log 25.$$

$$\begin{aligned} \log 25 &= \log \frac{100}{4} = \log 100 - \log (2 \times 2) \\ &= 2 - 2 \log 2 = 2 - .6020 = 1.3980. \end{aligned}$$

$$\therefore \log 25^{\frac{7}{3}} = \frac{7}{3} \times 1.3980 = 3.2620.$$

$$11. \log 15^{\frac{5}{6}} = \frac{5}{6} \log 15.$$

$$\log 15 = \log (3 \times 5) = \log 3 + \log 5.$$

$$\log 5 = \log \frac{10}{2} = 1 - \log 2 = .6990.$$

$$\therefore \log 15 = .4771 + .6990 = 1.1761.$$

$$\therefore \log 15^{\frac{5}{6}} = \frac{5}{6} \times 1.1761 = .9801.$$

$$12. \log \sqrt{7} = \frac{\log 7}{2} = \frac{.8451}{2} = .4225.$$

$$13. \log \sqrt[3]{3} = \frac{\log 3}{3} = \frac{.4771}{3} = .1590.$$

$$14. \log \sqrt[7]{2} = \frac{\log 2}{7} = \frac{.3010}{7} = .0430.$$

$$15. \log \sqrt[6]{5} = \frac{\log 5}{6}.$$

$$\log 5 = \log \frac{10}{2} = 1 - \log 2 = .6990.$$

$$\therefore \log \sqrt[6]{5} = \frac{.6990}{6} = .1165.$$

$$16. \log \sqrt[4]{35} = \frac{\log 35}{4}.$$

$$\log 35 = \log (5 \times 7) = \log 5 + \log 7.$$

$$\log 5 = \log \frac{10}{2} = 1 - \log 2 = .6990.$$

$$\therefore \log 35 = .6990 + .8451 = 1.5441.$$

$$\therefore \log \sqrt[4]{35} = \frac{1.5441}{4} = .3860.$$

$$17. \log \sqrt[9]{98} = \frac{\log 98}{9}.$$

$$\log 98 = \log (2 \times 7^2) = \log 2 + 2 \log 7 = .3010 + 1.6902 = 1.9912.$$

$$\therefore \log \sqrt[9]{98} = \frac{1.9912}{9} = .2212.$$

$$18. \log \sqrt[12]{126} = \frac{\log 126}{12}.$$

$$\begin{aligned} \log 126 &= \log (2 \times 3^2 \times 7) = \log 2 + 2 \log 3 + \log 7 \\ &= .3010 + .9542 + .8451 = 2.1003. \end{aligned}$$

$$\therefore \log \sqrt[12]{126} = \frac{2.1003}{12} = .1750.$$

$$\begin{aligned} 20. \log \left(\frac{10}{3} \right)^5 &= 5 \log \frac{10}{3} = 5 (\log 10 - \log 3) = 5 (1 - .4771) \\ &= 5 \times .5229 = 2.6145. \end{aligned}$$

$$\begin{aligned} 21. \log \frac{7^{\frac{3}{4}}}{5^{\frac{2}{3}}} &= \log 7^{\frac{3}{4}} - \log 5^{\frac{2}{3}} = \frac{3}{4} \log 7 - \frac{2}{3} \log 5 \\ &= .6338 - .4660 = .1678. \end{aligned}$$

$$\begin{aligned} 22. \log (3^{\frac{1}{6}} \times 2^{\frac{3}{5}}) &= \log 3^{\frac{1}{6}} + \log 2^{\frac{3}{5}} = \frac{1}{6} \log 3 + \frac{3}{5} \log 2 \\ &= .0795 + .1806 = .2601. \end{aligned}$$

$$\begin{aligned} 23. \log 3 \sqrt[4]{7} &= \log 3 + \log \sqrt[4]{7} = \log 3 + \frac{\log 7}{4} \\ &= .4771 + .2113 = .6884. \end{aligned}$$

$$24. \log \sqrt{\frac{7}{3}} = \frac{1}{2} \log \frac{7}{3} = \frac{\log 7 - \log 3}{2} = \frac{.8451 - .4771}{2} = .1840.$$

$$\begin{aligned} 25. \log \frac{\sqrt[3]{7}}{\sqrt[5]{2}} &= \log \sqrt[3]{7} - \log \sqrt[5]{2} = \frac{\log 7}{3} - \frac{\log 2}{5} \\ &= .2817 - .0602 = .2215. \end{aligned}$$

$$26. \log \sqrt[3]{\frac{28}{5}} = \frac{1}{3} \log \frac{28}{5} = \frac{\log 28 - \log 5}{3}.$$

$$\log 28 = \log (2^2 \times 7) = 2 \log 2 + \log 7 = .6020 + .8451 = 1.4471.$$

$$\therefore \log \sqrt[3]{\frac{28}{5}} = \frac{1.4471 - .6990}{3} = .2494.$$

$$27. \log \frac{\sqrt{42}}{10^{\frac{2}{3}}} = \log \sqrt{42} - \log 10^{\frac{2}{3}} = \frac{\log 42}{2} - \frac{2}{3} \log 10.$$

$$\begin{aligned} \log 42 &= \log (2 \times 3 \times 7) = \log 2 + \log 3 + \log 7 \\ &= .3010 + .4771 + .8451 = 1.6232. \end{aligned}$$

$$\therefore \log \frac{\sqrt{42}}{10^{\frac{2}{3}}} = .8116 - .6667 = .1449.$$

Art. 98.—Page 60.

2. $\log 18 = \log (2 \times 3^2) = \log 2 + 2 \log 3 = .3010 + .9542 = 1.2552$.
 $\therefore \log 1.8 = 0.2552$.
3. $\log 225 = \log (3^2 \times 5^2) = 2 \log 3 + 2 \log 5$
 $= .9542 + 1.3980 = 2.3522$.
 $\therefore \log 2.25 = 0.3522$.
4. $\log 196 = \log (2^2 \times 7^2) = 2 \log 2 + 2 \log 7$
 $= .6020 + 1.6902 = 2.2922$.
 $\therefore \log .196 = 2.2922 - 10$.
5. $\log 48 = \log (2^4 \times 3) = 4 \log 2 + \log 3 = 1.2040 + .4771 = 1.6811$.
 $\therefore \log .048 = 8.6811 - 10$.
6. $\log 384 = \log (2^7 \times 3) = 7 \log 2 + \log 3 = 2.1070 + .4771 = 2.5841$.
 $\therefore \log 38.4 = 1.5841$.
7. $\log 54 = \log (2 \times 3^3) = \log 2 + 3 \log 3 = .3010 + 1.4313 = 1.7323$.
 $\therefore \log .0054 = 7.7323 - 10$.
8. $\log 315 = \log (3^2 \times 5 \times 7) = 2 \log 3 + \log 5 + \log 7$
 $= .9542 + .6990 + .8451 = 2.4983$.
 $\therefore \log .000315 = 6.4983 - 10$.
9. $\log 735 = \log (3 \times 5 \times 7^2) = \log 3 + \log 5 + 2 \log 7$
 $= .4771 + .6990 + 1.6902 = 2.8663$.
 $\therefore \log 7350 = 3.8663$.
10. $\log 405 = \log (3^4 \times 5) = 4 \log 3 + \log 5$
 $= 1.9084 + .6990 = 2.6074$.
 $\therefore \log 4.05 = 0.6074$.
11. $\log 448 = \log (2^6 \times 7) = 6 \log 2 + \log 7$
 $= 1.8060 + .8451 = 2.6511$.
 $\therefore \log .448 = 9.6511 - 10$.
12. $\log 3024 = \log (2^4 \times 3^3 \times 7) = 4 \log 2 + 3 \log 3 + \log 7$
 $= 1.2040 + 1.4313 + .8451 = 3.4804$.
 $\therefore \log 302.4 = 2.4804$.
13. $\log 6174 = \log (2 \times 3^2 \times 7^3) = \log 2 + 2 \log 3 + 3 \log 7$
 $= .3010 + .9542 + 2.5353 = 3.7905$.
 $\therefore \log .06174 = 8.7905 - 10$.

$$14. \log (8.1)^7 = 7 \log 8.1.$$

$$\log 81 = \log 3^4 = 4 \log 3 = 1.9084.$$

$$\therefore \log 8.1 = 0.9084.$$

$$\therefore \log (8.1)^7 = 7 \times .9084 = 6.3588.$$

$$15. \log \sqrt[5]{9.6} = \frac{\log 9.6}{5}.$$

$$\begin{aligned} \log 96 &= \log (2^5 \times 3) = 5 \log 2 + \log 3 \\ &= 1.5050 + .4771 = 1.9821. \end{aligned}$$

$$\therefore \log 9.6 = 0.9821.$$

$$\therefore \log \sqrt[5]{9.6} = \frac{0.9821}{5} = .1964.$$

$$16. \log (22.4)^{\frac{1}{8}} = \frac{1}{8} \log 22.4.$$

$$\log 224 = \log (2^5 \times 7) = 5 \log 2 + \log 7 = 1.5050 + .8451 = 2.3501.$$

$$\therefore \log 22.4 = 1.3501.$$

$$\therefore \log (22.4)^{\frac{1}{8}} = \frac{1.3501}{8} = .1688.$$

Art. 105.—Pages 65 and 66.

$$1. \log (9.238 \times .9152) = \log 9.238 + \log .9152.$$

$$\log 9.238 = 0.9656$$

$$\log .9152 = 9.9615 - 10$$

$$\begin{array}{r} 0.9271 \\ \hline \end{array} = \log 8.454.$$

$$2. \log (130.36 \times .08237) = \log 130.36 + \log .08237.$$

$$\log 130.36 = 2.1151$$

$$\log .08237 = 8.9157 - 10$$

$$\begin{array}{r} 1.0308 \\ \hline \end{array} = \log 10.73.$$

$$3. \log (721.3 \times 3.0528) = \log 721.3 + \log 3.0528.$$

$$\log 721.3 = 2.8581$$

$$\log 3.0528 = 0.4847$$

$$\begin{array}{r} 3.3428 \\ \hline \end{array} = \log 2202.$$

Result, — 2202.

$$4. \log (4.3264 \times .050377) = \log 4.3264 + \log .050377.$$

$$\log 4.3264 = 0.6361$$

$$\log .050377 = 8.7022 - 10$$

$$\begin{array}{r} 9.3383 - 10 \\ \hline \end{array} = \log .2179.$$

$$\begin{aligned}
 5. \log (.27031 \times .042809) &= \log .27031 + \log .042809. \\
 \log .27031 &= 9.4319 - 10 \\
 \log .042809 &= 8.6315 - 10 \\
 \hline
 &8.0634 - 10 = \log .01157.
 \end{aligned}$$

$$\begin{aligned}
 6. \log (.063165 \times 11.134) &= \log .063165 + \log 11.134. \\
 \log .063165 &= 8.8005 - 10 \\
 \log 11.134 &= 1.0466 \\
 \hline
 &9.8471 - 10 = \log .7032. \\
 &\text{Result, } - .7032.
 \end{aligned}$$

$$\begin{aligned}
 7. \log \frac{401.8}{52.37} &= \log 401.8 - \log 52.37. \\
 \log 401.8 &= 2.6040 \\
 \log 52.37 &= 1.7191 \\
 \hline
 &0.8849 = \log 7.672.
 \end{aligned}$$

$$\begin{aligned}
 8. \log \frac{7.2321}{10.813} &= \log 7.2321 - \log 10.813. \\
 \log 7.2321 &= 0.8592 \\
 \log 10.813 &= 1.0339 \\
 \hline
 &9.8253 - 10 = \log .6688.
 \end{aligned}$$

$$\begin{aligned}
 9. \log \frac{.3384}{.08659} &= \log .3384 - \log .08659. \\
 \log .3384 &= 9.5294 - 10 \\
 \log .08659 &= 8.9374 - 10 \\
 \hline
 &0.5920 = \log 3.908. \\
 &\text{Result, } - 3.908.
 \end{aligned}$$

$$\begin{aligned}
 10. \log \frac{5.163}{.0051422} &= \log 5.163 - \log .0051422. \\
 \log 5.163 &= 0.9620 \\
 \log .0051422 &= 7.7112 - 10 \\
 \hline
 &3.2508 = \log 1782.
 \end{aligned}$$

$$\begin{aligned}
 11. \log \frac{22518}{64327} &= \log 22518 - \log 64327. \\
 \log 22518 &= 4.3525 \\
 \log 64327 &= 4.8084 \\
 \hline
 &9.5441 - 10 = \log .3500.
 \end{aligned}$$

$$12. \log \frac{.007514}{.015822} = \log .007514 - \log .015822.$$

$$\log .007514 = 7.8758 - 10$$

$$\log .015822 = 8.1993 - 10$$

$$\hline 9.6765 - 10 = \log .4748.$$

Result, — .4748.

$$13. \log \frac{3.3681}{12.853 \times .6349}$$

$$= \log 3.3681 + \text{colog } 12.853 + \text{colog } .6349.$$

$$\log 3.3681 = 0.5274$$

$$\text{colog } 12.853 = 8.8910 - 10$$

$$\text{colog } .6349 = 0.1973$$

$$\hline 9.6157 - 10 = \log .4127.$$

$$14. \log \frac{15.008 \times .0843}{.06376 \times 4.248}$$

$$= \log 15.008 + \log .0843 + \text{colog } .06376 + \text{colog } 4.248.$$

$$\log 15.008 = 1.1763$$

$$\log .0843 = 8.9258 - 10$$

$$\text{colog } .06376 = 1.1955$$

$$\text{colog } 4.248 = 9.3718 - 10$$

$$\hline 0.6694 = \log 4.671.$$

Result, — 4.671.

$$15. \log \frac{2563 \times .03442}{714.8 \times .511}$$

$$= \log 2563 + \log .03442 + \text{colog } 714.8 + \text{colog } .511.$$

$$\log 2563 = 3.4087$$

$$\log .03442 = 8.5368 - 10$$

$$\text{colog } 714.8 = 7.1458 - 10$$

$$\text{colog } .511 = 0.2916$$

$$\hline 9.3829 - 10 = \log .2415.$$

$$16. \log \frac{121.6 \times 9.025}{48.3 \times 3662 \times .0856} = \log 121.6$$

$$+ \log 9.025 + \text{colog } 48.3 + \text{colog } 3662 + \text{colog } .0856.$$

$$\log 121.6 = 2.0850$$

$$\log 9.025 = 0.9554$$

$$\text{colog } 48.3 = 8.3161 - 10$$

$$\text{colog } 3662 = 6.4363 - 10$$

$$\text{colog } .0856 = 1.0675$$

$$\hline 8.8603 - 10 = \log .0725.$$

Result, — .0725.

$$17. \log (23.86)^3 = 3 \times \log 23.86.$$

$$\log 23.86 = 1.3777$$

$$\underline{3}$$

$$4.1331 = \log 13587.$$

$$18. \log (.532)^8 = 8 \times \log .532.$$

$$\log .532 = 9.7259 - 10$$

$$\underline{8}$$

$$7.8072 - 10 = \log .006415.$$

$$19. \log (1.0246)^7 = 7 \times \log 1.0246.$$

$$\log 1.0246 = 0.0105$$

$$\underline{7}$$

$$0.0735 = \log 1.184.$$

$$\text{Result, } -1.184.$$

$$20. \log (.09323)^5 = 5 \times \log .09323.$$

$$\log .09323 = 8.9695 - 10$$

$$\underline{5}$$

$$4.8475 - 10 = \log .000007038.$$

$$21. \log 5^{\frac{2}{3}} = \frac{2}{3} \log 5.$$

$$\log 5 = 0.6990; \times \frac{2}{3} = 0.4660$$

$$= \log 2.924.$$

$$22. \log (.8)^{\frac{2}{5}} = \frac{2}{5} \log .8.$$

$$\log .8 = 9.9031 - 10$$

$$\underline{2}$$

$$5) 49.8062 - 50$$

$$9.9612 - 10 = \log .9146.$$

$$23. \log (3.16)^{\frac{4}{3}} = \frac{4}{3} \log 3.16.$$

$$\log 3.16 = 0.4997; \times \frac{4}{3} = 0.6663$$

$$= \log 4.638.$$

$$24. \log (.021)^{\frac{5}{2}} = \frac{5}{2} \log .021.$$

$$\log .021 = 8.3222 - 10$$

$$\underline{5}$$

$$2) 11.6110 - 20$$

$$5.8055 - 10 = \log .0000639.$$

$$25. \log \sqrt{2} = \frac{1}{2} \log 2.$$

$$\log 2 = 0.3010; \div 2 = 0.1505 \\ = \log 1.414.$$

$$26. \log \sqrt[4]{5} = \frac{1}{4} \log 5.$$

$$\log 5 = 0.6990; \div 4 = 0.1747 \\ = \log 1.495.$$

$$27. \log \sqrt[5]{3} = \frac{1}{5} \log 3.$$

$$\log 3 = 0.4771; \div 5 = 0.0954 \\ = \log 1.246. \\ \text{Result, } - 1.246.$$

$$28. \log \sqrt{.4294} = \frac{1}{2} \log .4294.$$

$$\log .4294 = 19.6329 - 20; \div 2 = 9.8164 - 10 \\ = \log .6553.$$

$$29. \log \sqrt[3]{.02305} = \frac{1}{3} \log .02305.$$

$$\log .02305 = 28.3626 - 30; \div 3 = 9.4542 - 10 \\ = \log .2846.$$

$$30. \log \sqrt[8]{1000} = \frac{1}{8} \log 1000.$$

$$\log 1000 = 3; \div 8 = 0.3750 = \log 2.372.$$

$$31. \log \sqrt[7]{.00951} = \frac{1}{7} \log .00951.$$

$$\log .00951 = 67.9782 - 70; \div 7 = 9.7112 - 10 \\ = \log .5142. \\ \text{Result, } - .5142.$$

$$32. \log \sqrt[5]{.0001011} = \frac{1}{5} \log .0001011.$$

$$\log .0001011 = 46.0047 - 50; \div 5 = 9.2009 - 10 \\ = \log .1588.$$

$$35. \log (2^{\frac{3}{2}} \times 3^{\frac{2}{3}}) = \frac{3}{2} \log 2 + \frac{2}{3} \log 3.$$

$$\log 2 = .3010; \times \frac{3}{2} = .4515$$

$$\log 3 = .4771; \times \frac{2}{3} = .3181$$

$$\underline{\hspace{1.5cm}} \\ .7696 = \log 5.883.$$

$$\begin{aligned}
 36. \log \frac{3^{\frac{5}{3}}}{4^{\frac{2}{3}}} &= \frac{5}{3} \log 3 - \frac{2}{3} \log 4. \\
 \log 3 &= .4771; \times \frac{5}{3} = .2982 \\
 \log 4 &= .6021; \times \frac{2}{3} = .4014 \\
 &\quad \underline{9.8968 - 10 = \log .7885.}
 \end{aligned}$$

$$\begin{aligned}
 37. \log \frac{5^{\frac{3}{7}}}{10^{\frac{2}{9}}} &= \frac{3}{7} \log 5 - \frac{2}{9} \log 10. \\
 \log 5 &= .6990; \times \frac{3}{7} = .2996 \\
 \log 10 &= 1; \times \frac{2}{9} = .2222 \\
 &\quad \underline{.0774 = \log 1.195.}
 \end{aligned}$$

$$\begin{aligned}
 38. \log \left(\frac{6}{7} \right)^{\frac{5}{2}} &= \frac{5}{2} (\log 6 - \log 7). \\
 \log 6 &= .7782 \\
 \log 7 &= .8451 \\
 &\quad \underline{9.9331 - 10} \\
 &\quad \quad \underline{5} \\
 &\quad 2) \underline{19.6655 - 20} \\
 &\quad \quad \underline{9.8327 - 10 = \log .6803.}
 \end{aligned}$$

$$\begin{aligned}
 39. \log \left(\frac{35}{113} \right)^{\frac{3}{8}} &= \frac{3}{8} (\log 35 - \log 113). \\
 \log 35 &= 1.5441 \\
 \log 113 &= 2.0531 \\
 &\quad \underline{9.4910 - 10} \\
 &\quad \quad \underline{3} \\
 &\quad 8) \underline{78.4730 - 80} \\
 &\quad \quad \underline{9.8091 - 10 = \log .6443.}
 \end{aligned}$$

$$\begin{aligned}
 40. \log \left(\frac{.08726}{.1321} \right)^{\frac{5}{3}} &= \frac{5}{3} (\log .08726 - \log .1321). \\
 \log .08726 &= 8.9408 - 10 \\
 \log .1321 &= 9.1209 - 10 \\
 &\quad \underline{9.8199 - 10} \\
 &\quad \quad \underline{5} \\
 &\quad 3) \underline{29.0995 - 30} \\
 &\quad \quad \underline{9.6998 - 10 = \log .5010.}
 \end{aligned}$$

$$41. \log \sqrt[8]{\frac{21}{13}} = \frac{1}{8} (\log 21 - \log 13).$$

$$\log 21 = 1.3222$$

$$\log 13 = 1.1139$$

$$\begin{array}{r} 8 \overline{) .2083} \end{array}$$

$$.0260 = \log 1.062.$$

$$42. \log \sqrt[9]{\frac{3}{7}} = \frac{1}{9} (\log 3 - \log 7).$$

$$\log 3 = .4771$$

$$\log 7 = .8451$$

$$\begin{array}{r} 9 \overline{) 89.6320 - 90} \end{array}$$

$$9.9591 - 10 = \log .9102.$$

$$\text{Result, } - .9102.$$

$$43. \log \left(\sqrt[5]{\frac{2}{3}} \div \sqrt[3]{\frac{3}{5}} \right)$$

$$= \frac{1}{5} (\log 2 - \log 3) - \frac{1}{3} (\log 3 - \log 5).$$

$$\log 2 = .3010$$

$$\log 3 = .4771$$

$$\log 3 = .4771$$

$$\log 5 = .6990$$

$$\begin{array}{r} 5 \overline{) 49.8239 - 50} \end{array}$$

$$9.9648 - 10$$

$$9.9260 - 10$$

$$.0388 = \log 1.093.$$

$$\begin{array}{r} 3 \overline{) 29.7781 - 30} \end{array}$$

$$9.9260 - 10$$

$$44. \log (\sqrt[8]{2} \times \sqrt[5]{3} \times \sqrt[7]{.01})$$

$$= \frac{1}{8} \log 2 + \frac{1}{5} \log 3 + \frac{1}{7} \log .01.$$

$$\log 2 = .3010; \quad \div 8 = .0376$$

$$\log 3 = .4771; \quad \div 5 = .0954$$

$$\log .01 = 68 - 70; \quad \div 7 = 9.7143 - 10$$

$$\begin{array}{r} 9.8473 - 10 = \log .7035. \end{array}$$

$$45. \log \sqrt[5]{\frac{3258}{49309}} = \frac{1}{5} (\log 3258 - \log 49309).$$

$$\log 3258 = 3.5129$$

$$\log 49309 = 4.6929$$

$$\begin{array}{r} 5 \overline{) 48.8200 - 50} \end{array}$$

$$9.7640 - 10 = \log .5807.$$

$$46. \log \left(\frac{31.63}{429} \right)^{\frac{3}{17}} = \frac{3}{17} (\log 31.63 - \log 429).$$

$$\log 31.63 = 1.5001$$

$$\log 429 = 2.6325$$

$$\underline{8.8676 - 10}$$

$$\underline{3}$$

$$17 \overline{) 166.6028 - 170}$$

$$9.8002 - 10 = \log .6313.$$

$$\text{Result, } -.6313.$$

$$47. \log \frac{100^{\frac{2}{3}}}{(.7325)^{\frac{3}{7}}} = \frac{2}{3} \log 100 - \frac{3}{7} \log .7325.$$

$$\log 100 = 2; \quad \times \frac{2}{3} = 1.3333$$

$$\log .7325 = 9.8648 - 10$$

$$\underline{3}$$

$$69.5944 - 70 \div 7 = \frac{9.9421 - 10}{1.3912 = \log 24.62.}$$

$$48. \log \frac{\sqrt[3]{.0001289}}{\sqrt[4]{.0008276}} = \frac{1}{3} \log .0001289 - \frac{1}{4} \log .0008276.$$

$$\log .0001289 = 26.1103 - 30; \div 3 = 8.7034 - 10$$

$$\log .0008276 = 36.9178 - 40; \div 4 = \frac{9.2294 - 10}{9.4740 - 10}$$

$$= \log .2979.$$

$$49. \log \frac{(.7469)^{\frac{5}{3}}}{(.2345)^{\frac{7}{2}}} = \frac{5}{3} \log .7469 - \frac{7}{2} \log .2345.$$

$$\log .7469 = 9.8732 - 10$$

$$\log .2345 = 9.3701 - 10$$

$$\underline{5}$$

$$3 \overline{) 29.3660 - 30}$$

$$9.7887 - 10$$

$$\underline{7.7953 - 10}$$

$$1.9934$$

$$= \log 98.50.$$

$$\underline{7}$$

$$2 \overline{) 15.5907 - 20}$$

$$7.7953 - 10$$

$$50. \log \frac{\sqrt[11]{.0073}}{(.68291)^{\frac{5}{2}}} = \frac{1}{11} \log .0073 - \frac{5}{2} \log .68291.$$

$$\log .0073 = 107.8633 - 110$$

$$\log .68291 = 9.8343 - 10$$

$$\text{Dividing by 11, } = 9.8058 - 10$$

$$\underline{9.5857 - 10}$$

$$.2201$$

$$= \log 1.660.$$

$$\underline{5}$$

$$2 \overline{) 19.1715 - 20}$$

$$9.5857 - 10$$

$$56. \log \frac{\sqrt[3]{.008193} \times (.06285)^{\frac{3}{2}}}{.98342}$$

$$= \frac{1}{3} \log .008193 + \frac{3}{2} \log .06285 + \text{colog} .98342.$$

$$\log .008193 = 27.9134 - 30; \div 3 = 9.3045 - 10$$

$$\log .06285 = 8.7983 - 10$$

$$\frac{3}{2}$$

$$\hline 16.3949 - 20; \div 2 = 8.1974 - 10$$

$$\text{colog} .98342 = 0.0072$$

$$\hline 7.5091 - 10$$

$$= \log .003229.$$

$$57. \log (\sqrt{.035} \times \sqrt[6]{.62667} \times \sqrt[3]{.0072103})$$

$$= \frac{1}{2} \log .035 + \frac{1}{6} \log .62667 + \frac{1}{3} \log .0072103.$$

$$\log .035 = 18.5441 - 20; \div 2 = 9.2720 - 10$$

$$\log .62667 = 59.7971 - 60; \div 6 = 9.9662 - 10$$

$$\log .0072103 = 27.8579 - 30; \div 3 = 9.2860 - 10$$

$$\hline 8.5242 - 10$$

$$= \log .03344.$$

CHAPTER VII.

Art. 110.—Pages 69 to 72.

- | | | |
|----|---|---|
| 1. | $a = c \sin A.$
$\log c = 1.0492$
$\log \sin A = \underline{9.8378}$
$\log a = 0.8870$
$\therefore a = 7.708.$ | $b = c \cos A.$
$\log c = 1.0492$
$\log \cos A = \underline{9.8606}$
$\log b = 0.9098$
$\therefore b = 8.124.$ |
| 2. | $b = a \tan B.$
$\log a = 2.8629$
$\log \tan B = \underline{0.4121}$
$\log b = 3.2750$
$\therefore b = 1883.$ | $c = \frac{a}{\cos B}.$
$\log a = 2.8629$
$\log \cos B = \underline{9.5576}$
$\log c = 3.3053$
$\therefore c = 2019.5.$ |
| 3. | $a = \frac{b}{\tan B}.$
$\log b = 1.6785$
$\log \tan B = \underline{0.2916}$
$\log a = 1.3869$
$\therefore a = 24.37.$ | $c = \frac{b}{\sin B}.$
$\log b = 1.6785$
$\log \sin B = \underline{9.9496}$
$\log c = 1.7289$
$\therefore c = 53.56.$ |
| 4. | $\sin A = \frac{a}{c}.$
$\log a = 9.7952$
$\log c = \underline{9.9590}$
$\log \sin A = 9.8362$
$\therefore A = 43^\circ 17.9'.$ | $b = \frac{a}{\tan A}.$
$\log a = 9.7952$
$\log \tan A = \underline{9.9742}$
$\log b = 9.8210$
$\therefore b = .6622.$ |
| 5. | $\tan A = \frac{a}{b}.$
$\log a = 0.6990$
$\log b = \underline{0.3010}$
$\log \tan A = 0.3980$
$\therefore A = 68^\circ 12.2'.$ | $c = \frac{a}{\sin A}.$
$\log a = 0.6990$
$\log \sin A = \underline{9.9678}$
$\log c = 0.7312$
$\therefore c = 5.385.$ |

6. $b = \frac{a}{\tan A}$ $c = \frac{a}{\sin A}$
 $\log a = 1.9212$ $\log a = 1.9212$
 $\log \tan A = 0.4912$ $\log \sin A = 9.9785$

 $\log b = 1.4300$ $\log c = 1.9427$
 $\therefore b = 26.91.$ $\therefore c = 87.64.$
7. $a = c \cos B$ $b = c \sin B.$
 $\log c = 8.4359$ $\log c = 8.4359$
 $\log \cos B = 9.9276$ $\log \sin B = 9.7262$

 $\log a = 8.3635$ $\log b = 8.1621$
 $\therefore a = .02309.$ $\therefore b = .01452.$
8. $\cos A = \frac{b}{c}$ $a = b \tan A.$
 $\log b = 0.4604$ $\log b = 0.4604$
 $\log c = 0.7084$ $\log \tan A = 0.1645$

 $\log \cos A = 9.7520$ $\log a = 0.6249$
 $\therefore A = 55^\circ 36.1'.$ $\therefore a = 4.216.$
9. $a = b \tan A.$ $c = \frac{b}{\cos A}$
 $\log b = 3.6281$ $\log b = 3.6281$
 $\log \tan A = 0.1179$ $\log \cos A = 9.7826$

 $\log a = 3.7460$ $\log c = 3.8455$
 $\therefore a = 5571.$ $\therefore c = 7007.$
10. $\tan A = \frac{a}{b}$ $c = \frac{a}{\sin A}$
 $\log a = 2.0043$ $\log a = 2.0043$
 $\log b = 2.0645$ $\log \sin A = 9.8173$

 $\log \tan A = 9.9398$ $\log c = 2.1870$
 $\therefore A = 41^\circ 2.4'.$ $\therefore c = 153.8.$
11. $b = \frac{a}{\tan A}$ $c = \frac{a}{\sin A}$
 $\log a = 2.1995$ $\log a = 2.1995$
 $\log \tan A = 9.9752$ $\log \sin A = 9.8368$

 $\log b = 2.2243$ $\log c = 2.3627$
 $\therefore b = 167.6.$ $\therefore c = 230.5.$

12. $a = c \sin A.$ $b = c \cos A.$
 $\log c = 1.5531$ $\log c = 1.5531$
 $\log \sin A = 9.9314$ $\log \cos A = 9.7162$

 $\log a = 1.4845$ $\log b = 1.2693$
 $\therefore a = 30.51.$ $\therefore b = 18.59.$
13. $\sin A = \frac{a}{c}.$ $b = \frac{a}{\tan A}.$
 $\log a = 2.3100$ $\log a = 2.3100$
 $\log c = 2.4398$ $\log \tan A = 0.0436$

 $\log \sin A = 9.8702$ $\log b = 2.2664$
 $\therefore A = 47^\circ 52.5'.$ $\therefore b = 184.7.$
14. $a = \frac{b}{\tan B}.$ $c = \frac{b}{\sin B}.$
 $\log b = 0.2158$ $\log b = 0.2158$
 $\log \tan B = 9.7614$ $\log \sin B = 9.6990$

 $\log a = 0.4544$ $\log c = 0.5168$
 $\therefore a = 2.847.$ $\therefore c = 3.287.$
15. $a = b \tan A.$ $c = \frac{b}{\cos A}.$
 $\log b = 1.1220$ $\log b = 1.1220$
 $\log \tan A = 9.6114$ $\log \cos A = 9.9665$

 $\log a = 0.7334$ $\log c = 1.1555$
 $\therefore a = 5.4125.$ $\therefore c = 14.306.$
16. $a = c \cos B.$ $b = c \sin B.$
 $\log c = 9.8611$ $\log c = 9.8611$
 $\log \cos B = 9.9922$ $\log \sin B = 9.2747$

 $\log a = 9.8533$ $\log b = 9.1358$
 $\therefore a = .7133.$ $\therefore b = .1367.$
17. $\tan A = \frac{a}{b}.$ $c = \frac{a}{\sin A}.$
 $\log a = 2.8051$ $\log a = 2.8051$
 $\log b = 2.7000$ $\log \sin A = 9.8957$

 $\log \tan A = 0.1051$ $\log c = 2.9094$
 $\therefore A = 51^\circ 51.9'.$ $\therefore c = 811.7.$

18. $b = \frac{a}{\tan A}$ $c = \frac{a}{\sin A}$
 $\log a = 2.3092$ $\log a = 2.3092$
 $\log \tan A = 0.6832$ $\log \sin A = 9.9908$

 $\log b = 1.6260$ $\log c = 2.3184$
 $\therefore b = 42.27.$ $\therefore c = 208.15.$
19. $\cos A = \frac{b}{c}$ $a = b \tan A$
 $\log b = 8.3974$ $\log b = 8.3974$
 $\log c = 8.6805$ $\log \tan A = 0.2143$

 $\log \cos A = 9.7169$ $\log a = 8.6117$
 $\therefore A = 58^\circ 35.7'.$ $\therefore a = .0409$
20. $b = a \tan B$ $c = \frac{a}{\cos B}$
 $\log a = 3.2731$ $\log a = 3.2731$
 $\log \tan B = 8.6085$ $\log \cos B = 9.9996$

 $\log b = 1.8816$ $\log c = 3.2735$
 $\therefore b = 76.13.$ $\therefore c = 1877.$
21. $\tan A = \frac{a}{b}$ $c = \frac{a}{\sin A}$
 $\log a = 1.3922$ $\log a = 1.3922$
 $\log b = 1.5188$ $\log \sin A = 9.7771$

 $\log \tan A = 9.8734$ $\log c = 1.6151$
 $\therefore A = 36^\circ 45.9'.$ $\therefore c = 41.22.$
22. $\cos A = \frac{b}{c}$ $a = b \tan A$
 $\log b = 0.1574$ $\log b = 0.1574$
 $\log c = 0.5397$ $\log \tan A = 0.3413$

 $\log \cos A = 9.6177$ $\log a = 0.4987$
 $\therefore A = 65^\circ 30'.$ $\therefore a = 3.153.$
23. $a = c \cos B$ $b = c \sin B$
 $\log c = 4.5706$ $\log c = 4.5706$
 $\log \cos B = 9.9975$ $\log \sin B = 9.0337$

 $\log a = 4.5681$ $\log b = 3.6043$
 $\therefore a = 36992.$ $\therefore b = 4021.$

24. $a = b \tan A.$

$$\begin{aligned}\log b &= 2.3011 \\ \log \tan A &= 0.3122 \\ \hline \log a &= 2.6133 \\ \therefore a &= 410.5.\end{aligned}$$

$$c = \frac{b}{\cos A}.$$

$$\begin{aligned}\log b &= 2.3011 \\ \log \cos A &= 9.6415 \\ \hline \log c &= 2.6596 \\ \therefore c &= 456.7.\end{aligned}$$

25. $\tan A = \frac{a}{b}.$

$$\begin{aligned}\log a &= 2.5316 \\ \log b &= 2.3649 \\ \hline \log \tan A &= 0.1667 \\ \therefore A &= 55^\circ 44.1' .\end{aligned}$$

$$c = \frac{a}{\sin A}.$$

$$\begin{aligned}\log a &= 2.5316 \\ \log \sin A &= 9.9172 \\ \hline \log c &= 2.6144 \\ \therefore c &= 411.5.\end{aligned}$$

26. $\sin A = \frac{a}{c}.$

$$\begin{aligned}\log a &= 0.2327 \\ \log c &= 0.3012 \\ \hline \log \sin A &= 9.9315 \\ \therefore A &= 58^\circ 40' .\end{aligned}$$

$$b = \frac{a}{\tan A}.$$

$$\begin{aligned}\log a &= 0.2327 \\ \log \tan A &= 0.2155 \\ \hline \log b &= 0.0172 \\ \therefore b &= 1.0405.\end{aligned}$$

27. $b = a \tan B.$

$$\begin{aligned}\log a &= 9.9144 \\ \log \tan B &= 9.5968 \\ \hline \log b &= 9.5112 \\ \therefore b &= .3245.\end{aligned}$$

$$c = \frac{a}{\cos B}.$$

$$\begin{aligned}\log a &= 9.9144 \\ \log \cos B &= 9.9685 \\ \hline \log c &= 9.9459 \\ \therefore c &= .8828.\end{aligned}$$

28. $a = c \sin A.$

$$\begin{aligned}\log c &= 2.4403 \\ \log \sin A &= 9.9828 \\ \hline \log a &= 2.4231 \\ \therefore a &= 264.9.\end{aligned}$$

$$b = c \cos A$$

$$\begin{aligned}\log c &= 2.4403 \\ \log \cos A &= 9.4402 \\ \hline \log b &= 1.8805 \\ \therefore b &= 75.95.\end{aligned}$$

29. $a = \frac{b}{\tan B}.$

$$\begin{aligned}\log b &= 2.0800 \\ \log \tan B &= 9.8329 \\ \hline \log a &= 2.2471 \\ \therefore a &= 176.64.\end{aligned}$$

$$c = \frac{b}{\sin B}.$$

$$\begin{aligned}\log b &= 2.0800 \\ \log \sin B &= 9.7503 \\ \hline \log c &= 2.3297 \\ \therefore c &= 213.65.\end{aligned}$$

$$\begin{array}{rcl}
 30. & \tan A = \frac{a}{b} & c = \frac{a}{\sin A} \\
 & \log a = 1.0046 & \log a = 1.0046 \\
 & \log b = 1.2381 & \log \sin A = 9.7027 \\
 & \log \tan A = 9.7665 & \log c = 1.3019 \\
 & \therefore A = 30^\circ 17.2'. & \therefore c = 20.04.
 \end{array}$$

$$\begin{array}{rcl}
 31. & c = 2b \cos A. & 34. \quad \cos A = \frac{\frac{1}{2}c}{a} \\
 & \log 2b = 1.8462 & \log \frac{1}{2}c = 1.7268 \\
 & \log \cos A = 9.5553 & \log a = 1.8989 \\
 & \log c = 1.4015 & \log \cos A = 9.8279 \\
 & \therefore c = 25.206. & \therefore A = 47^\circ 42.9'.
 \end{array}$$

$$\begin{array}{rcl}
 32. & a = \frac{\frac{1}{2}c}{\cos B} & 35. \quad a = \frac{\frac{1}{2}c}{\cos A} \\
 & \log \frac{1}{2}c = 0.1886 & \log \frac{1}{2}c = 9.4489 \\
 & \log \cos B = 9.9493 & \log \cos A = 9.5274 \\
 & \log a = 0.2393 & \log a = 9.9215 \\
 & \therefore a = 1.735. & \therefore a = .8346.
 \end{array}$$

$$\begin{array}{rcl}
 33. & c = 2b \sin \frac{1}{2}C. & 36. \quad a = \frac{\frac{1}{2}c}{\sin \frac{1}{2}C} \\
 & \log 2b = 3.6239 & \log \frac{1}{2}c = 1.6788 \\
 & \log \sin \frac{1}{2}C = 9.8116 & \log \sin \frac{1}{2}C = 9.9864 \\
 & \log c = 3.4355 & \log a = 1.6924 \\
 & \therefore c = 2725.6. & \therefore a = 49.25.
 \end{array}$$

37. Let O be the centre, and AB any side of the pentagon.

Join OA , and draw OC perpendicular to AB .

Then $AB = 2 OA \sin AOC = 24 \sin 36^\circ$.

$$\begin{array}{rcl}
 & \log 24 = 1.3802 & \\
 & \log \sin 36^\circ = 9.7692 & \\
 & \log AB = 1.1494 & \\
 & \therefore AB = 14.106. &
 \end{array}$$

38. Let A be the point of observation, B the top of the tower, and C its base.

Then $BC = AC \tan A = 100 \tan 38^\circ$.

$$\begin{array}{rcl}
 & \log 100 = 2.0000 & \\
 & \log \tan 38^\circ = 9.8928 & \\
 & \log BC = 1.8928 & \\
 & \therefore BC = 78.12. &
 \end{array}$$

39. Let B be the top and C the base of the tower, and A the extremity of its shadow.

Then

$$\begin{aligned}\tan A &= \frac{BC}{AC} = \frac{103.7}{167.3} \\ \log 103.7 &= 2.0157 \\ \log 167.3 &= 2.2235 \\ \log \tan A &= 9.7922 \\ \therefore A &= 31^\circ 47.1' .\end{aligned}$$

40. Let AB be the chord, and O the centre of the circle. Join OA and OB , and draw OC perpendicular to AB .

$$\begin{aligned}\text{Then} \quad \sin AOC &= \frac{AC}{OA} = \frac{513.5}{1634} \\ \log 513.5 &= 2.7105 \\ \log 1634 &= 3.2132 \\ \log \sin AOC &= 9.4973 \\ \therefore AOC &= 18^\circ 18.95' . \\ \therefore AOB &= 36^\circ 37.9' .\end{aligned}$$

41. Let A be the top of the mountain, B the remotest point visible, and O the centre of the earth.

Then in the right triangle OAB , $OA = 3956 + 1\frac{1}{4} = 3957.25$, and $OB = 3956$.

Hence

$$\begin{aligned}AB &= \sqrt{OA^2 - OB^2} \\ &= \sqrt{(OA + OB)(OA - OB)} \\ &= \sqrt{7913.25 \times 1.25} . \\ \log 7913.25 &= 3.8984 \\ \log 1.25 &= 0.0969 \\ &\quad \underline{2) 3.9953} \\ \log AB &= 1.9976 \\ \therefore AB &= 99.45 .\end{aligned}$$

42. Let AB and BC be consecutive sides of the pentagon. Join AC , and draw BD perpendicular to AC .

Then $AC = 2 AB \cos BAC = 14.056 \cos 36^\circ$.

$$\begin{aligned}\log 14.056 &= 1.1478 \\ \log \cos 36^\circ &= 9.9080 \\ \log AC &= 1.0558 \\ \therefore AC &= 11.371 .\end{aligned}$$

43. Let A denote the angle of elevation.

$$\begin{aligned}\text{Then} \quad \tan A &= \frac{238}{660} \\ \log 238 &= 2.3766 \\ \log 660 &= 2.8195 \\ \hline \log \tan A &= 9.5571 \\ \therefore A &= 19^\circ 50' .\end{aligned}$$

44. Let A be the position of the buoy, B the top of the light-house, and C its base.

$$\begin{aligned}\text{Then} \quad AC &= BC \cot BAC = 133 \cot 18^\circ 25' . \\ \log 133 &= 2.1239 \\ \log \cot 18^\circ 25' &= 0.4776 \\ \hline \log AC &= 2.6015 \\ \therefore AC &= 399.5 .\end{aligned}$$

45. Let H be the position of the headland, S the first point of observation, and S' the second.

Then in the right triangle HSS' , $SS' = 16.38$, and $\angle SHS' = 33^\circ$.

$$\text{Hence} \quad HS = 16.38 \cot 33^\circ ,$$

$$\text{and} \quad HS' = \frac{16.38}{\sin 33^\circ} .$$

$$\begin{aligned}\log 16.38 &= 1.2143 \\ \log \cot 33^\circ &= 0.1875 \\ \hline \log HS &= 1.4018 \\ \therefore HS &= 25.22 .\end{aligned}$$

$$\begin{aligned}\log 16.38 &= 1.2143 \\ \log \sin 33^\circ &= 9.7361 \\ \hline \log HS' &= 1.4782 \\ \therefore HS' &= 30.07 .\end{aligned}$$

46. Let AB be the chord, and O the centre of the circle. Join OA , and draw OC perpendicular to AB .

$$\begin{aligned}\text{Then} \quad OA &= \frac{AC}{\sin AOC} = \frac{20.68}{\sin 72^\circ 48.5'} . \\ \log 20.68 &= 1.3156 \\ \log \sin 72^\circ 48.5' &= 9.9801 \\ \hline \log OA &= 1.3355 \\ \therefore OA &= 21.65 .\end{aligned}$$

47. Let O be the centre, and AB any side of the octagon. Join OA , and draw OC perpendicular to AB .

Then $OC = AC \cot AOC = 6 \cot 22^\circ 30'$,

$$\text{and } OA = \frac{AC}{\sin AOC} = \frac{6}{\sin 22^\circ 30'}.$$

$$\log 6 = 0.7782$$

$$\log \cot 22^\circ 30' = 0.3828$$

$$\log OC = 1.1610$$

$$\therefore OC = 14.487.$$

$$\log 6 = 0.7782$$

$$\log \sin 22^\circ 30' = 9.5828$$

$$\log OA = 1.1954$$

$$\therefore OA = 15.682.$$

48. Let A be the position of the observer, B the top of the pole, and C its foot.

Then $AC = BC \cot BAC = 80 \cot 10^\circ$.

$$\log 80 = 1.9031$$

$$\log \cot 10^\circ = 0.7537$$

$$\log AC = 2.6568$$

$$\therefore AC = 453.7.$$

49. Let O be the centre, and AB any diagonal of the pentagon. Join OA , and draw OC perpendicular to AB .

Then $OA = \frac{AC}{\sin AOC} = \frac{16.415}{\sin 72^\circ}$.

$$\log 16.415 = 1.2152$$

$$\log \sin 72^\circ = 9.9782$$

$$\log OA = 1.2370$$

$$\therefore OA = 17.26.$$

50. Let B be the top, and C the foot of the tower, and A the extremity of the base line.

Then $BC = AC \tan BAC = 1000 \tan 21^\circ 16' 37''$.

$$\log 1000 = 3.0000$$

$$\log \tan 21^\circ 16' 37'' = 9.5904$$

$$\log BC = 2.5904$$

$$\therefore BC = 389.4.$$

51. Let AB be the chord, and O the centre of the circle.
Join OA , and draw OC perpendicular to AB .

Then $AB = 2 OA \sin AOC = 1446.58 \sin 17^\circ 36.5'.$

$$\log 1446.58 = 3.1604$$

$$\log \sin 17^\circ 36.5' = 9.4807$$

$$\log AB = \underline{2.6411}$$

$$\therefore AB = 437.6.$$

52. Let O be the centre, and AB any side of the hexagon.
Join OA , and draw OC perpendicular to AB .

Then $AB = 2 OC \tan AOC = 10 \tan 30^\circ.$

$$\log 10 = 1.0000$$

$$\log \tan 30^\circ = 9.7614$$

$$\log AB = \underline{0.7614}$$

$$\therefore AB = 5.773.$$

53. Let A be the position of the first boat, and B of the second; let C be the top of the light-house, and D its foot.

Then $AD = CD \cot CAD = 200 \cot 14^\circ,$

and $BD = CD \cot CBD = 200 \cot 32^\circ.$

$$\log 200 = 2.3010$$

$$\log \cot 14^\circ = 0.6032$$

$$\log AD = 2.9042$$

$$\therefore AD = 802.$$

$$\log 200 = 2.3010$$

$$\log \cot 32^\circ = 0.2042$$

$$\log BD = 2.5052$$

$$\therefore BD = 320.1.$$

$$\therefore AB = AD - BD = 481.9.$$

54. Let A be the position of the light-house, and B , C , and D , the positions of the ship at 7 A.M., 7.30 A.M., and 10 A.M., respectively.

Then $BC = AB \tan BAC = 10.32 \tan 18^\circ 13'.$

$$\log 10.32 = 1.0136$$

$$\log \tan 18^\circ 13' = 9.5174$$

$$\log BC = 0.5310$$

$$\therefore BC = 3.396.$$

Therefore the rate of the ship is 2×3.396 , or 6.792 miles an hour.

Again, $\tan BAD = \frac{BD}{AB} = \frac{20.376}{10.32}$
 $\log 20.376 = 1.3091$
 $\log 10.32 = 1.0136$
 $\log \tan BAD = 0.2955$
 $\therefore BAD = 63^\circ 8.4'$

Therefore the bearing of the light-house at 10 A.M. is $63^\circ 8.4'$ west of north.

Art. 112. — Page 74.

2. $2K = a^2 \cot A$
 $2 \log a = 2.6916$
 $\log \cot A = 0.4485$
 $\log 2K = 3.1401$
 $2K = 1380.6$
 $\therefore K = 690.3$

6. $4K = c^2 \sin 2A$
 $2 \log c = 4.5708$
 $\log \sin 2A = 9.9455$
 $\log 4K = 4.5163$
 $4K = 32831$
 $\therefore K = 8208$

3. $2K = a^2 \tan B$
 $2 \log a = 9.8290$
 $\log \tan B = 9.6510$
 $\log 2K = 9.4800$
 $2K = .302$
 $\therefore K = .151$

7. $2K = b^2 \tan A$
 $2 \log b = 7.4332$
 $\log \tan A = 0.2190$
 $\log 2K = 7.6522$
 $2K = .00449$
 $\therefore K = .002245$

4. $2K = ab$
 $\log a = 2.1741$
 $\log b = 1.8824$
 $\log 2K = 4.0565$
 $2K = 11389$
 $\therefore K = 5695$

8. $2K = a\sqrt{(c+a)(c-a)}$
 $\log a = 9.9694$
 $\frac{1}{2} \log (c+a) = 0.2851$
 $\frac{1}{2} \log (c-a) = 0.1341$
 $\log 2K = 0.3886$
 $2K = 2.447$
 $\therefore K = 1.223$

5. $2K = b\sqrt{(c+b)(c-b)}$
 $\log b = 9.4851$
 $\frac{1}{2} \log (c+b) = 9.9924$
 $\frac{1}{2} \log (c-b) = 9.7748$
 $\log 2K = 9.2523$
 $2K = .17876$
 $\therefore K = .08938$

9. $4K = c^2 \sin 2B$
 $2 \log c = 2.8718$
 $\log \sin 2B = 9.7604$
 $\log 4K = 2.6322$
 $4K = 428.7$
 $\therefore K = 107.2$

10. $2K = b^2 \cot B$
 $2 \log b = 9.0574$
 $\log \cot B = 0.2508$
 $\log 2K = 9.3082$
 $2K = .2033$
 $\therefore K = .1017$

CHAPTER IX.

Art. 121.—Page 84.

2. $C = 180^\circ - 115^\circ 10' = 64^\circ 50'.$

$$b = a \sin B \csc A.$$

$$\log a = 1.0000$$

$$\log \sin B = 9.9890$$

$$\log \csc A = 0.2107$$

$$\log b = 1.1997$$

$$\therefore b = 15.837.$$

$$c = a \sin C \csc A.$$

$$\log a = 1.0000$$

$$\log \sin C = 9.9567$$

$$\log \csc A = 0.2107$$

$$\log c = 1.1674$$

$$\therefore c = 14.703.$$

3. $A = 180^\circ - 154^\circ = 26^\circ.$

$$a = b \sin A \csc B.$$

$$\log b = 9.9051$$

$$\log \sin A = 9.6418$$

$$\log \csc B = 0.1015$$

$$\log a = 9.6484$$

$$\therefore a = .445.$$

$$c = b \sin C \csc B.$$

$$\log b = 9.9051$$

$$\log \sin C = 9.9909$$

$$\log \csc B = 0.1015$$

$$\log c = 9.9975$$

$$\therefore c = .9942.$$

4. $C = 180^\circ - 80^\circ 35' = 99^\circ 25'.$

$$a = c \sin A \csc C.$$

$$\log c = 8.5051$$

$$\log \sin A = 9.7706$$

$$\log \csc C = 0.0059$$

$$\log a = 8.2816$$

$$\therefore a = .01913.$$

$$b = c \sin B \csc C.$$

$$\log c = 8.5051$$

$$\log \sin B = 9.8453$$

$$\log \csc C = 0.0059$$

$$\log b = 8.3563$$

$$\therefore b = .02272.$$

5. $B = 180^\circ - 120^\circ 55' = 59^\circ 5'.$

$$a = b \sin A \csc B.$$

$$\log b = 1.4625$$

$$\log \sin A = 9.9996$$

$$\log \csc B = 0.0666$$

$$\log a = 1.5287$$

$$\therefore a = 33.78.$$

$$c = b \sin C \csc B.$$

$$\log b = 1.4625$$

$$\log \sin C = 9.7390$$

$$\log \csc B = 0.0666$$

$$\log c = 1.2681$$

$$\therefore c = 18.54.$$

6. $A = 180^\circ - 139^\circ 23' = 40^\circ 37'.$

$$\begin{aligned} b &= a \sin B \csc A. \\ \log a &= 0.7340 \\ \log \sin B &= 9.9954 \\ \log \csc A &= \underline{0.1865} \\ \log b &= 0.9159 \\ \therefore b &= 8.24. \end{aligned}$$

$$\begin{aligned} c &= a \sin C \csc A. \\ \log a &= 0.7340 \\ \log \sin C &= 9.8170 \\ \log \csc A &= \underline{0.1865} \\ \log c &= 0.7375 \\ \therefore c &= 5.464. \end{aligned}$$

7. $B = 180^\circ - 158^\circ 54' = 21^\circ 6'.$

$$\begin{aligned} a &= c \sin A \csc C. \\ \log c &= 8.2068 \\ \log \sin A &= 9.7613 \\ \log \csc C &= \underline{0.0796} \\ \log a &= 8.0477 \\ \therefore a &= .011162. \end{aligned}$$

$$\begin{aligned} b &= c \sin B \csc C. \\ \log c &= 8.2068 \\ \log \sin B &= 9.5563 \\ \log \csc C &= \underline{0.0796} \\ \log b &= 7.8427 \\ \therefore b &= .006962. \end{aligned}$$

8. $B = 180^\circ - 114^\circ 28' = 65^\circ 32'.$

$$\begin{aligned} b &= a \sin B \csc A. \\ \log a &= 2.6021 \\ \log \sin B &= 9.9591 \\ \log \csc A &= \underline{0.0895} \\ \log b &= 2.6507 \\ \therefore b &= 447.4. \end{aligned}$$

$$\begin{aligned} c &= a \sin C \csc A. \\ \log a &= 2.6021 \\ \log \sin C &= 9.9375 \\ \log \csc A &= \underline{0.0895} \\ \log c &= 2.6291 \\ \therefore c &= 425.7. \end{aligned}$$

9. $C = 180^\circ - 125^\circ 13' = 54^\circ 47'.$

$$\begin{aligned} a &= b \sin A \csc B. \\ \log b &= 2.4973 \\ \log \sin A &= 9.9652 \\ \log \csc B &= \underline{0.0723} \\ \log a &= 2.5348 \\ \therefore a &= 342.6. \end{aligned}$$

$$\begin{aligned} c &= b \sin C \csc B. \\ \log b &= 2.4973 \\ \log \sin C &= 9.9122 \\ \log \csc B &= \underline{0.0723} \\ \log c &= 2.4818 \\ \therefore c &= 303.3. \end{aligned}$$

10. $A = 180^\circ - 75^\circ 28' 18'' = 104^\circ 31' 42''.$

$$\begin{aligned} a &= c \sin A \csc C. \\ \log c &= 0.8954 \\ \log \sin A &= 9.9858 \\ \log \csc C &= \underline{0.1628} \\ \log a &= 1.0440 \\ \therefore a &= 11.067. \end{aligned}$$

$$\begin{aligned} b &= c \sin B \csc C. \\ \log c &= 0.8954 \\ \log \sin B &= 9.7248 \\ \log \csc C &= \underline{0.1628} \\ \log b &= 0.7830 \\ \therefore b &= 6.067. \end{aligned}$$

Art. 122. — Page 85.

$$2. \tan \frac{1}{2}(A - C) = \frac{a - c}{a + c} \tan \frac{1}{2}(A + C).$$

$$b = a \sin B \csc A.$$

$$a - c = 12 \quad \log = 1.0792$$

$$\log a = 1.4314$$

$$a + c = 42 \quad \text{colog} = 8.3768$$

$$\log \sin B = 9.8569$$

$$\frac{1}{2}(A + C) = 67^\circ \quad \log \tan = 0.3721$$

$$\log \csc A = 0.0080$$

$$\log \tan \frac{1}{2}(A - C) = 9.8281$$

$$\log b = 1.2963$$

$$\therefore \frac{1}{2}(A - C) = 33^\circ 56.7'.$$

$$\therefore b = 19.78.$$

$$\therefore A = \frac{1}{2}(A + C) + \frac{1}{2}(A - C) = 100^\circ 56.7',$$

$$\text{and } C = \frac{1}{2}(A + C) - \frac{1}{2}(A - C) = 33^\circ 3.3'.$$

$$3. \tan \frac{1}{2}(A - B) = \frac{a - b}{a + b} \tan \frac{1}{2}(A + B).$$

$$c = a \sin C \csc A.$$

$$a - b = 139 \quad \log = 2.1430$$

$$\log a = 2.6866$$

$$a + b = 833 \quad \text{colog} = 7.0794$$

$$\log \sin C = 9.8941$$

$$\frac{1}{2}(A + B) = 64^\circ 12' \quad \log \tan = 0.3157$$

$$\log \csc A = 0.0030$$

$$\log \tan \frac{1}{2}(A - B) = 9.5381$$

$$\log c = 2.5837$$

$$\therefore \frac{1}{2}(A - B) = 19^\circ 2.7'.$$

$$\therefore c = 383.5.$$

$$\therefore A = \frac{1}{2}(A + B) + \frac{1}{2}(A - B) = 83^\circ 14.7',$$

$$\text{and } B = \frac{1}{2}(A + B) - \frac{1}{2}(A - B) = 45^\circ 9.3'.$$

$$4. \tan \frac{1}{2}(C - B) = \frac{c - b}{c + b} \tan \frac{1}{2}(C + B).$$

$$a = b \sin A \csc B.$$

$$c - b = 1.265 \quad \log = 0.1021$$

$$\log b = 0.3621$$

$$c + b = 5.869 \quad \text{colog} = 9.2315$$

$$\log \sin A = 9.9459$$

$$\frac{1}{2}(C + B) = 59^\circ \quad \log \tan = 0.2212$$

$$\log \csc B = 0.1987$$

$$\log \tan \frac{1}{2}(C - B) = 9.5548$$

$$\log a = 0.5067$$

$$\therefore \frac{1}{2}(C - B) = 19^\circ 44.2'.$$

$$\therefore a = 3.211.$$

$$\therefore C = \frac{1}{2}(C + B) + \frac{1}{2}(C - B) = 78^\circ 44.2',$$

$$\text{and } B = \frac{1}{2}(C + B) - \frac{1}{2}(C - B) = 39^\circ 15.8'.$$

$$5. \tan \frac{1}{2}(B - A) = \frac{b - a}{b + a} \tan \frac{1}{2}(B + A).$$

$$c = a \sin C \csc A.$$

$$b - a = .063 \quad \log = 8.7993$$

$$\log a = 9.4771$$

$$b + a = .663 \quad \text{colog} = 0.1785$$

$$\log \sin C = 9.9137$$

$$\frac{1}{2}(B + A) = 27^\circ 32' \quad \log \tan = 9.7171$$

$$\log \csc A = 0.3790$$

$$\log \tan \frac{1}{2}(B - A) = 8.6949$$

$$\log c = 9.7698$$

$$\therefore \frac{1}{2}(B - A) = 2^\circ 50.2'.$$

$$\therefore c = .5886.$$

$$\therefore B = \frac{1}{2}(B + A) + \frac{1}{2}(B - A) = 30^\circ 22.2',$$

$$\text{and } A = \frac{1}{2}(B + A) - \frac{1}{2}(B - A) = 24^\circ 41.8'.$$

$$6. \tan \frac{1}{2}(B - C) = \frac{b - c}{b + c} \tan \frac{1}{2}(B + C).$$

$$b - c = 835.8 \quad \log = 2.9221$$

$$b + c = 1548.4 \quad \text{colog} = 6.8101$$

$$\frac{1}{2}(B + C) = 76^\circ 52' \quad \log \tan = \underline{0.6320}$$

$$\log \tan \frac{1}{2}(B - C) = 0.3642$$

$$\therefore \frac{1}{2}(B - C) = 66^\circ 37.1'.$$

$$\therefore B = \frac{1}{2}(B + C) + \frac{1}{2}(B - C) = 143^\circ 29.1',$$

$$\text{and } C = \frac{1}{2}(B + C) - \frac{1}{2}(B - C) = 10^\circ 14.9'.$$

$$a = b \sin A \csc B.$$

$$\log b = 3.0763$$

$$\log \sin A = 9.6460$$

$$\log \csc B = \underline{0.2254}$$

$$\log a = 2.9477$$

$$\therefore a = 886.6.$$

$$7. \tan \frac{1}{2}(C - A) = \frac{c - a}{c + a} \tan \frac{1}{2}(C + A).$$

$$c - a = 4.039 \quad \log = 0.6063$$

$$c + a = 18.839 \quad \text{colog} = 8.7249$$

$$\frac{1}{2}(C + A) = 48^\circ 47' \quad \log \tan = \underline{0.0575}$$

$$\log \tan \frac{1}{2}(C - A) = 9.3887$$

$$\therefore \frac{1}{2}(C - A) = 13^\circ 45.1'.$$

$$\therefore C = \frac{1}{2}(C + A) + \frac{1}{2}(C - A) = 62^\circ 32.1',$$

$$\text{and } A = \frac{1}{2}(C + A) - \frac{1}{2}(C - A) = 35^\circ 1.9'.$$

$$b = a \sin B \csc A.$$

$$\log a = 0.8692$$

$$\log \sin B = 9.9962$$

$$\log \csc A = \underline{0.2411}$$

$$\log b = 1.1065$$

$$\therefore b = 12.78.$$

$$8. \tan \frac{1}{2}(A - B) = \frac{a - b}{a + b} \tan \frac{1}{2}(A + B).$$

$$a - b = 11.66 \quad \log = 1.0667$$

$$a + b = 94.88 \quad \text{colog} = 8.0228$$

$$\frac{1}{2}(A + B) = 50^\circ 43.5' \quad \log \tan = \underline{0.0874}$$

$$\log \tan \frac{1}{2}(A - B) = 9.1769$$

$$\therefore \frac{1}{2}(A - B) = 8^\circ 32.8'.$$

$$\therefore A = \frac{1}{2}(A + B) + \frac{1}{2}(A - B) = 59^\circ 16.3',$$

$$\text{and } B = \frac{1}{2}(A + B) - \frac{1}{2}(A - B) = 42^\circ 10.7'.$$

$$c = a \sin C \csc A.$$

$$\log a = 1.7265$$

$$\log \sin C = 9.9913$$

$$\log \csc A = \underline{0.0657}$$

$$\log c = 1.7835$$

$$\therefore c = 60.74.$$

$$9. \tan \frac{1}{2}(C - B) = \frac{c - b}{c + b} \tan \frac{1}{2}(C + B).$$

$$c - b = .02424 \quad \log = 8.3845$$

$$c + b = .0776 \quad \text{colog} = 1.1101$$

$$\frac{1}{2}(C + B) = 32^\circ 6.5' \quad \log \tan = \underline{9.7976}$$

$$\log \tan \frac{1}{2}(C - B) = 9.2922$$

$$\therefore \frac{1}{2}(C - B) = 11^\circ 5.3'.$$

$$\therefore C = \frac{1}{2}(C + B) + \frac{1}{2}(C - B) = 43^\circ 11.8',$$

$$\text{and } B = \frac{1}{2}(C + B) - \frac{1}{2}(C - B) = 21^\circ 1.2'.$$

$$a = b \sin A \csc B.$$

$$\log b = 8.4262$$

$$\log \sin A = 9.9545$$

$$\log \csc B = \underline{0.4453}$$

$$\log a = 8.8260$$

$$\therefore a = .06699.$$

$$\begin{array}{ll}
 10. \tan \frac{1}{2}(C-A) = \frac{c-a}{c+a} \tan \frac{1}{2}(C+A). & b = a \sin B \csc A. \\
 c-a = 16.56 & \log = 1.2191 \\
 c+a = 119.32 & \csc \log = 7.9233 \\
 \frac{1}{2}(C+A) = 50^\circ 23' 43'' & \log \tan = 0.0823 \\
 \log \tan \frac{1}{2}(C-A) = 9.2247 & \log b = 1.8873 \\
 \therefore \frac{1}{2}(C-A) = 9^\circ 31.4' = 9^\circ 31' 24''. & \therefore b = 77.14. \\
 \therefore C = \frac{1}{2}(C+A) + \frac{1}{2}(C-A) = 59^\circ 55' 7'', \\
 \text{and } A = \frac{1}{2}(C+A) - \frac{1}{2}(C-A) = 40^\circ 52' 19''.
 \end{array}$$

Art. 123. — Page 88.

3. Here $s = 4.5$, $s-a = 2.5$, $s-b = 1.5$, $s-c = .5$.

$$\begin{array}{ll}
 \log(s-a) = 0.3979 & \log r = 9.8099 \\
 \log(s-b) = 0.1761 & \log(s-b) = 0.1761 \\
 \log(s-c) = 9.6990 & \log \tan \frac{1}{2}B = 9.6338 \\
 \csc \log s = 9.3468 & \frac{1}{2}B = 23^\circ 17.1'. \\
 2) 9.6198 & \therefore B = 46^\circ 34.2'. \\
 \log r = 9.8099 & \log r = 9.8099 \\
 \log r = 9.8099 & \log(s-c) = 9.6990 \\
 \log(s-a) = 0.3979 & \log \tan \frac{1}{2}C = 0.1109 \\
 \log \tan \frac{1}{2}A = 9.4120 & \frac{1}{2}C = 52^\circ 14.2'. \\
 \frac{1}{2}A = 14^\circ 28.7'. & \therefore C = 104^\circ 28.4'. \\
 \therefore A = 28^\circ 57.4'.
 \end{array}$$

Check, $A + B + C = 180^\circ$.

4. Here $s = 8.5$, $s-a = 4.5$, $s-b = 1.5$, $s-c = 2.5$.

$$\begin{array}{ll}
 \log(s-a) = 0.6532 & \log r = 0.1489 \\
 \log(s-b) = 0.1761 & \log(s-b) = 0.1761 \\
 \log(s-c) = 0.3979 & \log \tan \frac{1}{2}B = 9.9728 \\
 \csc \log s = 9.0706 & \frac{1}{2}B = 43^\circ 12.4'. \\
 2) 0.2978 & \therefore B = 86^\circ 24.8'. \\
 \log r = 0.1489 & \log r = 0.1489 \\
 \log(s-a) = 0.6532 & \log(s-c) = 0.3979 \\
 \log \tan \frac{1}{2}A = 9.4957 & \log \tan \frac{1}{2}C = 9.7510 \\
 \frac{1}{2}A = 17^\circ 23.2'. & \frac{1}{2}C = 29^\circ 24.5'. \\
 \therefore A = 34^\circ 46.4'. & \therefore C = 58^\circ 49'.
 \end{array}$$

Check, $A + B + C = 180^\circ 0.2$.

5. Here $s = 7.4$, $s - a = 1.8$, $s - b = 3.1$, $s - c = 2.5$.

$$\log(s - a) = 0.2553$$

$$\log(s - b) = 0.4914$$

$$\log(s - c) = 0.3979$$

$$\text{colog } s = 9.1308$$

$$2) 0.2754$$

$$\log r = 0.1377$$

$$\log(s - a) = 0.2553$$

$$\log \tan \frac{1}{2} A = 9.8824$$

$$\frac{1}{2} A = 37^\circ 20'.$$

$$\therefore A = 74^\circ 40'.$$

$$\log r = 0.1377$$

$$\log(s - b) = 0.4914$$

$$\log \tan \frac{1}{2} B = 9.6463$$

$$\frac{1}{2} B = 23^\circ 53.2'.$$

$$\therefore B = 47^\circ 46.4'.$$

$$\log r = 0.1377$$

$$\log(s - c) = 0.3979$$

$$\log \tan \frac{1}{2} C = 9.7398$$

$$\frac{1}{2} C = 28^\circ 46.7'.$$

$$\therefore C = 57^\circ 33.4'.$$

$$\text{Check, } A + B + C = 179^\circ 59.8'.$$

6. Here $s = .344$, $s - a = .114$, $s - b = .084$, $s - c = .146$.

$$\log(s - a) = 9.0569$$

$$\log(s - b) = 8.9243$$

$$\log(s - c) = 9.1644$$

$$\text{colog } s = 0.4634$$

$$2) 7.6090$$

$$\log r = 8.8045$$

$$\log(s - a) = 9.0569$$

$$\log \tan \frac{1}{2} A = 9.7476$$

$$\frac{1}{2} A = 29^\circ 13'.$$

$$\therefore A = 58^\circ 26'.$$

$$\log r = 8.8045$$

$$\log(s - b) = 8.9243$$

$$\log \tan \frac{1}{2} B = 9.8802$$

$$\frac{1}{2} B = 37^\circ 11.9'.$$

$$\therefore B = 74^\circ 23.8'.$$

$$\log r = 8.8045$$

$$\log(s - c) = 9.1644$$

$$\log \tan \frac{1}{2} C = 9.6401$$

$$\frac{1}{2} C = 23^\circ 35.3'.$$

$$\therefore C = 47^\circ 10.6'.$$

$$\text{Check, } A + B + C = 180^\circ 0.4'.$$

7. Here $s = 120.2$, $s - a = 40.9$, $s - b = 26$, $s - c = 53.3$.

$$\log(s - a) = 1.6117$$

$$\log(s - b) = 1.4150$$

$$\log(s - c) = 1.7267$$

$$\text{colog } s = 7.9201$$

$$2) 2.6735$$

$$\log r = 1.3367$$

$$\log(s - a) = 1.6117$$

$$\log \tan \frac{1}{2} A = 9.7250$$

$$\frac{1}{2} A = 27^\circ 57.7'.$$

$$\therefore A = 55^\circ 55.4'.$$

$$\log r = 1.3367$$

$$\log(s - b) = 1.4150$$

$$\log \tan \frac{1}{2} B = 9.9217$$

$$\frac{1}{2} B = 39^\circ 51.9'.$$

$$\therefore B = 79^\circ 43.8'.$$

$$\log r = 1.3367$$

$$\log(s - c) = 1.7267$$

$$\log \tan \frac{1}{2} C = 9.6100$$

$$\frac{1}{2} C = 22^\circ 10'.$$

$$\therefore C = 44^\circ 20'.$$

$$\text{Check, } A + B + C = 179^\circ 59.2'.$$

8. Here
- $s = 542$
- ,
- $s - a = 221$
- ,
- $s - b = 181$
- ,
- $s - c = 140$
- .

$$\begin{aligned}\log(s - a) &= 2.3444 \\ \log(s - b) &= 2.2577 \\ \log(s - c) &= 2.1461 \\ \text{colog } s &= 7.2660\end{aligned}$$

$$\begin{aligned}&2) 4.0142 \\ &\log r = 2.0071 \\ \log(s - a) &= 2.3444 \\ \log \tan \frac{1}{2} A &= 9.6627 \\ \frac{1}{2} A &= 24^\circ 42.1'. \\ \therefore A &= 49^\circ 24.2'.\end{aligned}$$

$$\begin{aligned}\log r &= 2.0071 \\ \log(s - b) &= 2.2577 \\ \log \tan \frac{1}{2} B &= 9.7494 \\ \frac{1}{2} B &= 29^\circ 19'. \\ \therefore B &= 58^\circ 38'. \\ \log r &= 2.0071 \\ \log(s - c) &= 2.1461 \\ \log \tan \frac{1}{2} C &= 9.8610 \\ \frac{1}{2} C &= 35^\circ 58.9'. \\ \therefore C &= 71^\circ 57.8'.\end{aligned}$$

Check, $A + B + C = 180^\circ$.

9. Here
- $s = .936$
- ,
- $s - a = .295$
- ,
- $s - b = .407$
- ,
- $s - c = .234$
- .

$$\begin{aligned}\log(s - a) &= 9.4698 \\ \log(s - b) &= 9.6096 \\ \log(s - c) &= 9.3692 \\ \text{colog } s &= 0.0287\end{aligned}$$

$$\begin{aligned}&2) 8.4773 \\ &\log r = 9.2386 \\ \log(s - a) &= 9.4698 \\ \log \tan \frac{1}{2} A &= 9.7688 \\ \frac{1}{2} A &= 30^\circ 25.4'. \\ \therefore A &= 60^\circ 50.8'.\end{aligned}$$

$$\begin{aligned}\log r &= 9.2386 \\ \log(s - b) &= 9.6096 \\ \log \tan \frac{1}{2} B &= 9.6290 \\ \frac{1}{2} B &= 23^\circ 3.1'. \\ \therefore B &= 46^\circ 6.2'. \\ \log r &= 9.2386 \\ \log(s - c) &= 9.3692 \\ \log \tan \frac{1}{2} C &= 9.8694 \\ \frac{1}{2} C &= 36^\circ 30.8'. \\ \therefore C &= 73^\circ 1.6'.\end{aligned}$$

Check, $A + B + C = 179^\circ 58.6'$.

10. Here
- $s = 6.989$
- ,
- $s - a = 3.97$
- ,
- $s - b = .258$
- ,
- $s - c = 2.761$
- .

$$\begin{aligned}\log(s - a) &= 0.5988 \\ \log(s - b) &= 9.4116 \\ \log(s - c) &= 0.4411 \\ \text{colog } s &= 9.1556\end{aligned}$$

$$\begin{aligned}&2) 9.6071 \\ &\log r = 9.8035 \\ \log(s - a) &= 0.5988 \\ \log \tan \frac{1}{2} A &= 9.2047 \\ \frac{1}{2} A &= 9^\circ 6.2'. \\ \therefore A &= 18^\circ 12.4'.\end{aligned}$$

$$\begin{aligned}\log r &= 9.8035 \\ \log(s - b) &= 9.4116 \\ \log \tan \frac{1}{2} B &= 0.3919 \\ \frac{1}{2} B &= 67^\circ 55.3'. \\ \therefore B &= 135^\circ 50.6'. \\ \log r &= 9.8035 \\ \log(s - c) &= 0.4411 \\ \log \tan \frac{1}{2} C &= 9.3624 \\ \frac{1}{2} C &= 12^\circ 58.3'. \\ \therefore C &= 25^\circ 56.6'.\end{aligned}$$

Check, $A + B + C = 179^\circ 59.6'$.

Art. 127. — Pages 91 and 92.

6. Since b is $< a$, there is but one solution, corresponding to the acute value of B .

$$\sin B = \frac{b \sin A}{a}$$

$$c = a \sin C \csc A.$$

$$\log b = 0.5551$$

$$\log a = 0.7059$$

$$\operatorname{colog} a = 9.2941$$

$$\log \sin C = 9.9884$$

$$\log \sin A = 9.9530$$

$$\log \csc A = 0.0470$$

$$\log \sin B = 9.8022$$

$$\log c = 0.7413$$

$$\therefore B = 39^\circ 21.3',$$

$$\therefore c = 5.511.$$

and

$$C = 180^\circ - 103^\circ 11.3' = 76^\circ 48.7'.$$

7. Since b is $> c$, and C is acute, there will be two solutions, one solution, or no solution, according as $\log \sin B$ is negative, zero, or positive.

$$\sin B = \frac{b \sin C}{c}$$

$$a_1 = b \sin A_1 \csc B.$$

$$a_2 = b \sin A_2 \csc B.$$

$$\log b = 1.8739$$

$$\log b = 1.8739$$

$$\log b = 1.8739$$

$$\operatorname{colog} c = 8.2062$$

$$\log \sin A_1 = 9.9408$$

$$\log \sin A_2 = 9.0316$$

$$\log \sin C = 9.6615$$

$$\log \csc B = 0.2584$$

$$\log \csc B = 0.2584$$

$$\log \sin B = 9.7416$$

$$\log a_1 = 2.0731$$

$$\log a_2 = 1.1639$$

$$\therefore B_1 = 33^\circ 28.4',$$

$$\therefore a_1 = 118.33.$$

$$\therefore a_2 = 14.58.$$

and

$$B_2 = 146^\circ 31.6'.$$

$$\therefore A_1 = 180^\circ - 60^\circ 46.4' = 119^\circ 13.6',$$

and

$$A_2 = 180^\circ - 173^\circ 49.6' = 6^\circ 10.4'.$$

8. Since c is $< b$, there is but one solution, corresponding to the acute value of C .

$$\sin C = \frac{c \sin B}{b}$$

$$a = b \sin A \csc B.$$

$$\log c = 9.2971$$

$$\log b = 9.3687$$

$$\operatorname{colog} b = 0.6313$$

$$\log \sin A = 9.4825$$

$$\log \sin B = 9.9757$$

$$\log \csc B = 0.0243$$

$$\log \sin C = 9.9041$$

$$\log a = 8.8755$$

$$\therefore C = 53^\circ 18.9',$$

$$\therefore a = .07508.$$

and

$$A = 180^\circ - 162^\circ 18.9' = 17^\circ 41.1'.$$

9. Since a is $< c$, there is but one solution, corresponding to the acute value of A .

$$\sin A = \frac{a \sin C}{c}.$$

$$b = a \sin B \csc A.$$

$$\log a = 0.0294$$

$$\log a = 0.0294$$

$$\text{colog } c = 9.7670$$

$$\log \sin B = 9.8916$$

$$\log \sin C = 9.7228$$

$$\log \csc A = 0.4808$$

$$\log \sin A = 9.5192$$

$$\log b = 0.4018$$

$$\therefore A = 19^\circ 18.1',$$

$$\therefore b = 2.522.$$

and $B = 180^\circ - 51^\circ 11.1' = 128^\circ 48.9'.$

$$10. \sin A = \frac{a \sin B}{b}.$$

$$c_1 = a \sin C_1 \csc A.$$

$$c_2 = a \sin C_2 \csc A.$$

$$\log a = 9.2704$$

$$\log a = 9.2704$$

$$\log a = 9.2704$$

$$\text{colog } b = 0.7696$$

$$\log \sin C_1 = 9.7795$$

$$\log \sin C_2 = 9.4314$$

$$\log \sin B = 9.9524$$

$$\log \csc A = 0.0076$$

$$\log \csc A = 0.0076$$

$$\log \sin A = 9.9924$$

$$\log c_1 = 9.0575$$

$$\log c_2 = 8.7094$$

$$\therefore A_1 = 79^\circ 20',$$

$$\therefore c_1 = .11416.$$

$$\therefore c_2 = .05121$$

and $A_2 = 100^\circ 40'.$

$$\therefore C_1 = 180^\circ - 143^\circ = 37^\circ,$$

and $C_2 = 180^\circ - 164^\circ 20' = 15^\circ 40'.$

11. Since c is $> a$, and A is obtuse, the triangle is impossible.

12. Since b is $< c$, there is but one solution, corresponding to the acute value of B .

$$\sin B = \frac{b \sin C}{c}.$$

$$a = b \sin A \csc B.$$

$$\log b = 1.7016$$

$$\log b = 1.7016$$

$$\text{colog } c = 8.1752$$

$$\log \sin A = 9.9232$$

$$\log \sin C = 9.7340$$

$$\log \csc B = 0.3892$$

$$\log \sin B = 9.6108$$

$$\log a = 2.0140$$

$$\therefore B = 24^\circ 5.4',$$

$$\therefore a = 103.3.$$

and $A = 180^\circ - 56^\circ 54.4' = 123^\circ 5.6'.$

$$13. \sin C = \frac{c \sin A}{a}.$$

$$b = a \tan B.$$

$$\log c = 1.0000$$

$$\log a = 0.9373$$

$$\text{colog } a = 9.0627$$

$$\log \tan B = 9.7623$$

$$\log \sin A = 9.9373$$

$$\log b = 0.6996$$

$$0.0000$$

$$\therefore b = 5.007.$$

$$\therefore C = 90^\circ,$$

and $B = 90^\circ - 59^\circ 57' = 30^\circ 3'.$

$$14. \quad \sin C = \frac{c \sin B}{b}. \quad a_1 = b \sin A_1 \csc B. \quad a_2 = b \sin A_2 \csc B.$$

$$\log c = 0.8351$$

$$\log b = 0.7127$$

$$\log b = 0.7127$$

$$\text{colog } b = 9.2873$$

$$\log \sin A_1 = 9.9695$$

$$\log \sin A_2 = 9.5939$$

$$\log \sin B = 9.8422$$

$$\log \csc B = 0.1578$$

$$\log \csc B = 0.1578$$

$$\log \sin C = 9.9646$$

$$\log a_1 = 0.8400$$

$$\log a_2 = 0.4644$$

$$\therefore C_1 = 67^\circ 10',$$

$$\therefore a_1 = 6.918.$$

$$\therefore a_2 = 2.913.$$

$$\text{and } C_2 = 112^\circ 50'.$$

$$\therefore A_1 = 180^\circ - 111^\circ 13' = 68^\circ 47',$$

$$\text{and } A_2 = 180^\circ - 156^\circ 53' = 23^\circ 7'.$$

15. Since a is $< b$, there is only one solution, corresponding to the acute value of A .

$$\sin A = \frac{a \sin B}{b}.$$

$$c = a \sin C \csc A.$$

$$\log a = 2.3315$$

$$\log a = 2.3315$$

$$\text{colog } b = 7.5455$$

$$\log \sin C = 9.6825$$

$$\log \sin B = 9.9863$$

$$\log \csc A = 0.1367$$

$$\log \sin A = 9.8633$$

$$\log c = 2.1507$$

$$\therefore A = 46^\circ 53.3',$$

$$\therefore c = 141.48.$$

$$\text{and } C = 180^\circ - 151^\circ 13.3' = 28^\circ 46.7'.$$

$$16. \quad \sin B = \frac{b \sin C}{c}.$$

$$\log b = 3.4870$$

$$\text{colog } c = 6.9126$$

$$\log \sin C = 9.9179$$

$$\log \sin B = 0.3175$$

Since $\log \sin B$ is positive, the triangle is impossible.

17. Since c is $< a$, there is only one solution, corresponding to the acute value of C .

$$\sin C = \frac{c \sin A}{a}.$$

$$b = a \sin B \csc A.$$

$$\log c = 9.7086$$

$$\log a = 9.8511$$

$$\text{colog } a = 0.1489$$

$$\log \sin B = 9.9363$$

$$\log \sin A = 9.7606$$

$$\log \csc A = 0.2394$$

$$\log \sin C = 9.6181$$

$$\log b = 0.0268$$

$$\therefore C = 24^\circ 31.4',$$

$$\therefore b = 1.0637.$$

$$\text{and } B = 180^\circ - 59^\circ 42.4' = 120^\circ 17.6'.$$

$$18. \quad \sin B = \frac{b \sin A}{a} \qquad c = b \sin C.$$

$$\log b = 2.2206$$

$$\log b = 2.2206$$

$$\text{colog } a = 7.9712$$

$$\log \sin C = 9.8843$$

$$\log \sin A = 9.8082$$

$$\log c = 2.1049$$

$$\log \sin B = 0.0000$$

$$\therefore c = 127.32.$$

$$\therefore B = 90^\circ,$$

$$\text{and} \quad C = 90^\circ - 40^\circ 0' 21'' = 49^\circ 59' 39''.$$

$$19. \quad \sin A = \frac{a \sin C}{c} \quad b_1 = a \sin B_1 \csc A. \quad b_2 = a \sin B_2 \csc A.$$

$$\log a = 9.5073$$

$$\log a = 9.5073$$

$$\log a = 9.5073$$

$$\text{colog } c = 0.5673$$

$$\log \sin B_1 = 9.9255$$

$$\log \sin B_2 = 9.4853$$

$$\log \sin C = 9.8989$$

$$\log \csc A = 0.0265$$

$$\log \csc A = 0.0265$$

$$\log \sin A = 9.9735$$

$$\log b_1 = 9.4593$$

$$\log b_2 = 9.0191$$

$$\therefore A_1 = 70^\circ 12',$$

$$\therefore b_1 = .2879.$$

$$\therefore b_2 = .1045.$$

$$\text{and} \quad A_2 = 109^\circ 48'.$$

$$\therefore B_1 = 180^\circ - 122^\circ 36' = 57^\circ 24',$$

$$\text{and} \quad B_2 = 180^\circ - 162^\circ 12' = 17^\circ 48'.$$

20. Since c is $< b$, there is only one solution, corresponding to the acute value of C .

$$\sin C = \frac{c \sin B}{b} \qquad a = b \sin A \csc B.$$

$$\log c = 2.7828$$

$$\log b = 2.9092$$

$$\text{colog } b = 7.0908$$

$$\log \sin A = 9.4596$$

$$\log \sin B = 9.9075$$

$$\log \csc B = 0.0925$$

$$\log \sin C = 9.7811$$

$$\log a = 2.4613$$

$$\therefore C = 37^\circ 10',$$

$$\therefore a = 289.3.$$

$$\text{and} \quad A = 180^\circ - 163^\circ 15' 20'' = 16^\circ 44' 40''.$$

Art. 128.—Page 93.

$$2. \quad 2K = ac \sin B.$$

$$3. \quad \text{Here} \quad s = 9,$$

$$\log a = 1.5798$$

$$s - a = 4,$$

$$\log c = 1.7868$$

$$s - b = 2,$$

$$\log \sin B = 9.9670$$

$$\text{and} \quad s - c = 3.$$

$$\log 2K = 3.3336$$

$$K = \sqrt{s(s-a)(s-b)(s-c)}.$$

$$2K = 2155.7$$

$$\therefore K = 1077.9.$$

$$\log s = 0.9542$$

$$\log (s - a) = 0.6021$$

$$\log (s - b) = 0.3010$$

$$\log (s - c) = 0.4771$$

$$2) \underline{2.3344}$$

$$\log K = 1.1672$$

$$\therefore K = 14.697.$$

$$4. \quad 2K = b^2 \sin C \sin A \csc B.$$

$$C = 180^\circ - 106^\circ 23' = 73^\circ 37'.$$

$$2 \log b = 0.6320$$

$$\log \sin C = 9.9820$$

$$\log \sin A = 9.9730$$

$$\log \csc B = 0.2268$$

$$\log 2K = 0.8138$$

$$2K = 6.513$$

$$\therefore K = 3.257.$$

$$5. \quad 2K = bc \sin A.$$

$$\log b = 2.0649$$

$$\log c = 2.0000$$

$$\log \sin A = 9.9449$$

$$\log 2K = 4.0098$$

$$2K = 10229$$

$$\therefore K = 5114.$$

$$6. \text{ Here } s = 120,$$

$$s - a = 41,$$

$$s - b = 26,$$

$$\text{and } s - c = 53.$$

$$K = \sqrt{s(s-a)(s-b)(s-c)}.$$

$$\log s = 2.0792$$

$$\log (s - a) = 1.6128$$

$$\log (s - b) = 1.4150$$

$$\log (s - c) = 1.7243$$

$$2) \underline{6.8313}$$

$$\log K = 3.4156$$

$$\therefore K = 2604.$$

$$7. \quad 2K = a^2 \sin B \sin C \csc A.$$

$$C = 180^\circ - 67^\circ 8' = 112^\circ 52'.$$

$$2 \log a = 0.9892$$

$$\log \sin B = 9.3822$$

$$\log \sin C = 9.9645$$

$$\log \csc A = 0.0966$$

$$\log 2K = 0.4325$$

$$2K = 2.707$$

$$\therefore K = 1.353.$$

$$8. \quad 2K = b^2 \sin C \sin A \csc B.$$

$$B = 180^\circ - 117^\circ 13' = 62^\circ 47'.$$

$$2 \log b = 9.2850$$

$$\log \sin C = 9.8132$$

$$\log \sin A = 9.9880$$

$$\log \csc B = 0.0510$$

$$\log 2K = 9.1372$$

$$2K = .13716$$

$$\therefore K = .06858.$$

$$9. \text{ Here } s = 34,$$

$$s - a = 10.9,$$

$$s - b = 14.3,$$

$$\text{and } s - c = 8.8.$$

$$K = \sqrt{s(s-a)(s-b)(s-c)}.$$

$$\log s = 1.5315$$

$$\log (s - a) = 1.0374$$

$$\log (s - b) = 1.1553$$

$$\log (s - c) = 0.9445$$

$$2) \underline{4.6687}$$

$$\log K = 2.3343$$

$$\therefore K = 215.9.$$

$$10. \quad 2K = ac \sin B.$$

$$\log a = 9.5089$$

$$\log c = 9.9582$$

$$\log \sin B = 9.9387$$

$$\log 2K = 9.4058$$

$$2K = .2546$$

$$\therefore K = .1273.$$

$$\begin{aligned}
 11. \quad 2K &= c^2 \sin A \sin B \csc C. \\
 A &= 180^\circ - 131^\circ 49' = 48^\circ 11'. \\
 2 \log c &= 3.8088 \\
 \log \sin A &= 9.8723 \\
 \log \sin B &= 9.9932 \\
 \log \csc C &= 0.2791 \\
 \hline
 \log 2K &= 3.9534 \\
 2K &= 8982 \\
 \therefore K &= 4491.
 \end{aligned}$$

$$\begin{aligned}
 12. \quad 2K &= ab \sin C. \\
 \log a &= 8.0072 \\
 \log b &= 8.2607 \\
 \log \sin C &= 9.2924 \\
 \hline
 \log 2K &= 5.5603 \\
 2K &= .00003633 \\
 \therefore K &= .00001817.
 \end{aligned}$$

$$\begin{aligned}
 13. \quad \text{Here} \quad s &= 8.04, \\
 s - a &= 2.22, \\
 s - b &= 2.04, \\
 \text{and} \quad s - c &= 3.78. \\
 K &= \sqrt{s(s-a)(s-b)(s-c)}. \\
 \log s &= 0.9053 \\
 \log (s-a) &= 0.3464 \\
 \log (s-b) &= 0.3096 \\
 \log (s-c) &= 0.5775
 \end{aligned}$$

$$\begin{aligned}
 &2) 2.1388 \\
 \log K &= 1.0694 \\
 \therefore K &= 11.732.
 \end{aligned}$$

Art. 129.— Pages 93 and 94.

1. Let A be the first point of observation, and B the second; and let C be the top of the tower, and D its base.

$$\begin{aligned}
 \text{Then} \quad \frac{AC}{AB} &= \frac{\sin ABC}{\sin ACB}, \\
 \text{or,} \quad AC &= AB \sin ABC \csc ACB \\
 &= 100 \sin 35^\circ 16' \csc 17^\circ 23'. \\
 \log 100 &= 2.0000 \\
 \log \sin 35^\circ 16' &= 9.7615 \\
 \log \csc 17^\circ 23' &= 0.5247 \\
 \hline
 \therefore \log AC &= 2.2862
 \end{aligned}$$

$$\begin{aligned}
 \text{Then} \quad CD &= AC \sin CAD = AC \sin 52^\circ 39', \\
 \text{and} \quad AD &= AC \cos CAD = AC \cos 52^\circ 39'. \\
 \log AC &= 2.2862 \\
 \log \sin 52^\circ 39' &= 9.9003 \\
 \hline
 \log CD &= 2.1865 \\
 \therefore CD &= 153.64. \\
 \log AC &= 2.2862 \\
 \log \cos 52^\circ 39' &= 9.7830 \\
 \hline
 \log AD &= 2.0692 \\
 \therefore AD &= 117.27. \\
 \therefore BD &= AB + AD = 217.27.
 \end{aligned}$$

2. Denoting the sides opposite the angles A , B , and C of the triangle ABC by a , b , and c , respectively, we have

$$K = \sqrt{s(s-a)(s-b)(s-c)}.$$

Here $s = 351$, $s - a = 115$, $s - b = 40$, and $s - c = 196$.

$$\begin{aligned} \log s &= 2.5453 \\ \log (s - a) &= 2.0607 \\ \log (s - b) &= 1.6021 \\ \log (s - c) &= 2.2923 \\ \hline 2) 8.5004 \\ \log K &= 4.2502 \\ \therefore K &= 17792. \end{aligned}$$

Denoting the sides opposite the angles A , C , and D of the triangle ACD by a , c , and d , respectively, we have

$$K = \sqrt{s(s-a)(s-c)(s-d)}.$$

Here $s = 334$, $s - a = 82$, $s - c = 229$, and $s - d = 23$.

$$\begin{aligned} \log s &= 2.5237 \\ \log (s - a) &= 1.9138 \\ \log (s - c) &= 2.3598 \\ \log (s - d) &= 1.3617 \\ \hline 2) 8.1590 \\ \log K &= 4.0795 \\ \therefore K &= 12008. \end{aligned}$$

Therefore $\text{area } ABCD = 17792 + 12008 = 29800$.

3. Let A be the position of the first post, and B of the second, and let C be the top of the bluff, and D its foot.

$$\text{Then} \quad \frac{AC}{AB} = \frac{\sin ABC}{\sin ACB}.$$

$$\begin{aligned} \text{Whence,} \quad CD &= AC \sin CAD \\ &= \frac{AB \sin ABC \sin CAD}{\sin ACB} \\ &= \frac{1000 \sin 9^\circ 33' \sin 27^\circ 40'}{\sin 18^\circ 7'}. \end{aligned}$$

$$\begin{aligned} \log 1000 &= 3.0000 \\ \log \sin 9^\circ 33' &= 9.2198 \\ \log \sin 27^\circ 40' &= 9.6668 \\ \log \csc 18^\circ 7' &= 0.5073 \\ \hline \log CD &= 2.3939 \\ \therefore CD &= 247.7. \end{aligned}$$

4. Let A be the starting-point, and B and C the positions of the first and second yachts, respectively, at the end of 40 minutes.

Then in the triangle ABC , we have

$$AB = 6.96, \quad AC = 5.14, \quad \text{and } \angle A = 45^\circ.$$

$$\tan \frac{1}{2}(C - B) = \frac{AB - AC}{AB + AC} \tan \frac{1}{2}(C + B).$$

$$AB - AC = 1.82 \qquad \log = 0.2601$$

$$AB + AC = 12.1 \qquad \text{colog} = 8.9172$$

$$\frac{1}{2}(C + B) = 67^\circ 30' \qquad \log \tan = 0.3828$$

$$\log \tan \frac{1}{2}(C - B) = 9.5601$$

$$\therefore \frac{1}{2}(C - B) = 19^\circ 57.5',$$

$$\text{and } B = \frac{1}{2}(C + B) - \frac{1}{2}(C - B) = 47^\circ 32.5'.$$

$$BC = AC \sin A \csc B.$$

$$\log AC = 0.7110$$

$$\log \sin A = 9.8495$$

$$\log \csc B = 0.1321$$

$$\log BC = 0.6926$$

$$\therefore BC = 4.927.$$

5. Let A be the position of the lighthouse, and B and C the first and second positions of the ship.

Then in the triangle ABC , we have

$$BC = 14, \quad \angle B = 105^\circ, \quad \text{and } \angle C = 30^\circ.$$

$$\text{Whence, } \angle A = 180^\circ - 135^\circ = 45^\circ.$$

$$AB = BC \sin C \csc A.$$

$$\log BC = 1.1461$$

$$\log \sin C = 9.6990$$

$$\log \csc A = 0.1505$$

$$\log AB = 0.9956$$

$$\therefore AB = 9.9.$$

$$AC = BC \sin B \csc A.$$

$$\log BC = 1.1461$$

$$\log \sin B = 9.9849$$

$$\log \csc A = 0.1505$$

$$\log AC = 1.2815$$

$$\therefore AC = 19.122.$$

6.

$$AB = BC \sin C \csc A.$$

$$A = 180^\circ - 158^\circ 23' = 21^\circ 37'.$$

$$\log BC = 2.3187$$

$$\log \sin C = 9.7218$$

$$\log \csc A = \underline{0.4337}$$

$$\log AB = 2.4742$$

$$\therefore AB = 298.$$

7. Let A be the point of observation, B the top of the flag-pole, C its foot, and D the base of the tower.

Then,
$$\frac{AC}{BC} = \frac{\sin ABC}{\sin BAC},$$

or,

$$AC = BC \sin ABC \csc BAC \\ = 40 \sin 51^\circ 7' \csc 18^\circ 35'.$$

$$\log 40 = 1.6021$$

$$\log \sin 51^\circ 7' = 9.8912$$

$$\log \csc 18^\circ 35' = \underline{0.4967}$$

$$\therefore \log AC = 1.9900$$

$$AD = AC \cos CAD = AC \cos 20^\circ 18'.$$

$$\log AC = 1.9900$$

$$\log \cos 20^\circ 18' = \underline{9.9722}$$

$$\log AD = 1.9622$$

$$\therefore AD = 91.66.$$

$$CD = AC \sin CAD = AC \sin 20^\circ 18'.$$

$$\log AC = 1.9900$$

$$\log \sin 20^\circ 18' = \underline{9.5402}$$

$$\log CD = 1.5302$$

$$\therefore CD = 33.9.$$

8.

$$BC = BD \sin BDC \csc BCD$$

$$= BD \sin 60^\circ \csc 20^\circ.$$

But,

$$BD = AD \sin BAD \csc ABD$$

$$= 500 \sin 60^\circ \csc 80^\circ.$$

Whence,

$$BC = 500 \sin^2 60^\circ \csc 20^\circ \csc 80^\circ.$$

$$\log 500 = 2.6990$$

$$2 \log \sin 60^\circ = 9.8750$$

$$\log \csc 20^\circ = 0.4659$$

$$\log \csc 80^\circ = \underline{0.0066}$$

$$\log BC = 3.0465$$

$$\therefore BC = 1113.1.$$

9. $AC = CD \sin ADC \csc CAD.$

$$\log CD = 2.1761$$

$$\log \sin ADC = 9.6990$$

$$\log \csc CAD = 0.0866$$

$$\log AC = 1.9617$$

$$\therefore AC = 91.56.$$

$$BC = CD \sin BDC \csc CBD.$$

$$\log CD = 2.1761$$

$$\log \sin BDC = 9.9968$$

$$\log \csc CBD = 0.3430$$

$$\log BC = 2.5159$$

$$\therefore BC = 328.$$

$$\tan \frac{1}{2}(BAC - ABC) = \frac{BC - AC}{BC + AC} \tan \frac{1}{2}(BAC + ABC)$$

$$BC - AC = 236.44$$

$$\log = 2.3737$$

$$BC + AC = 419.56$$

$$\text{colog} = 7.3772$$

$$\frac{1}{2}(BAC + ABC) = 77^\circ 30'$$

$$\log \tan = 0.6542$$

$$\log \tan \frac{1}{2}(BAC - ABC) = 0.4051$$

$$\therefore \frac{1}{2}(BAC - ABC) = 68^\circ 31.4',$$

and $BAC = \frac{1}{2}(BAC + ABC) + \frac{1}{2}(BAC - ABC) = 146^\circ 1.4'.$

$$AB = BC \sin ACB \csc BAC.$$

$$\log BC = 2.5159$$

$$\log \sin ACB = 9.6259$$

$$\log \csc BAC = 0.2527$$

$$\log AB = 2.3945$$

$$\therefore AB = 248.$$

10. Denoting the sides opposite the angles A , B , and C of the triangle ABC by a , b , and c , respectively, we have

$$K = \sqrt{s(s-a)(s-b)(s-c)}.$$

Here,

$$s = 87.5,$$

$$s - a = 24.5,$$

$$s - b = 12.5,$$

and

$$s - c = 50.5.$$

$$\log s = 1.9420$$

$$\log (s - a) = 1.3892$$

$$\log (s - b) = 1.0969$$

$$\log (s - c) = 1.7033$$

$$2) 6.1314$$

$$\log K = 3.0657$$

$$\therefore K = 1163.2.$$

$$\cos \frac{1}{2} BAC = \sqrt{\frac{s(s-a)}{bc}}.$$

$$\log s = 1.9420$$

$$\log (s-a) = 1.3892$$

$$\text{colog } b = 8.1249$$

$$\text{colog } c = 8.4318$$

$$2)9.8879$$

$$\log \cos \frac{1}{2} BAC = 9.9439$$

$$\frac{1}{2} BAC = 28^{\circ} 30'$$

$$\therefore BAC = 57^{\circ}.$$

Denoting the sides opposite the angles A , B , and D of the triangle ABD by a , b , and d , respectively, we have

$$\cos \frac{1}{2} BAD = \sqrt{\frac{s(s-a)}{bd}}.$$

Here,
and

$$s = 49.5,$$

$$s-a = 7.5.$$

$$\log s = 1.6946$$

$$\log (s-a) = 0.8751$$

$$\text{colog } b = 8.6990$$

$$\text{colog } d = 8.4318$$

$$2)9.7005$$

$$\log \cos \frac{1}{2} BAD = 9.8502$$

$$\frac{1}{2} BAD = 44^{\circ} 54.2'$$

$$\therefore BAD = 89^{\circ} 48.4'.$$

$$\therefore CAD = BAD - BAC = 32^{\circ} 48.4'.$$

$$2 \text{ area } ACD = AC \cdot AD \cdot \sin CAD.$$

$$\log AC = 1.8751$$

$$\log AD = 1.3010$$

$$\log \sin CAD = 9.7339$$

$$\log (2 \text{ area } ACD) = 2.9100$$

$$2 \text{ area } ACD = 812.8$$

$$\therefore \text{area } ACD = 406.4.$$

$$\therefore \text{area } ABCD = 1163.2 + 406.4 = 1569.6.$$

CHAPTER XI.

Art. 153.—Page 112.

$$\begin{aligned}
 5. \quad \sin A &= \frac{\sin a}{\sin c} \\
 \log \sin a &= 9.5543 \\
 \log \sin c &= 9.8311 \\
 \log \sin A &= 9.7232 \\
 180^\circ - A &= 31^\circ 55'. \\
 \therefore A &= 148^\circ 5'.
 \end{aligned}$$

$$\begin{aligned}
 \cos B &= \frac{\tan a}{\tan c} \\
 \log \tan a &= 9.5842 \\
 \log \tan c &= 9.9646 \\
 \log \cos B &= 9.6196 \\
 \therefore B &= 65^\circ 23.2'.
 \end{aligned}$$

$$\begin{aligned}
 \cos b &= \frac{\cos c}{\cos a} \\
 \log \cos c &= 9.8665 \\
 \log \cos a &= 9.9702 \\
 \log \cos b &= 9.8963 \\
 \therefore b &= 38^\circ 2'.
 \end{aligned}$$

Check.

$$\begin{aligned}
 \sin A &= \frac{\cos B}{\cos b} \\
 \log \cos B &= 9.6196 \\
 \log \cos b &= 9.8963 \\
 \log \sin A &= 9.7233
 \end{aligned}$$

$$\begin{aligned}
 6. \quad \cos a &= \frac{\cos A}{\sin B} \\
 \log \cos A &= 9.8050 \\
 \log \sin B &= 9.9252 \\
 \log \cos a &= 9.8798 \\
 \therefore a &= 40^\circ 41.8'.
 \end{aligned}$$

$$\begin{aligned}
 \cos b &= \frac{\cos B}{\sin A} \\
 \log \cos B &= 9.7322 \\
 \log \sin A &= 9.8864 \\
 \log \cos b &= 9.8458 \\
 180^\circ - b &= 45^\circ 29.2'. \\
 \therefore b &= 134^\circ 30.8'.
 \end{aligned}$$

$$\begin{aligned}
 \cos c &= \cot A \cot B. \\
 \log \cot A &= 9.9187 \\
 \log \cot B &= 9.8070 \\
 \log \cos c &= 9.7257 \\
 180^\circ - c &= 57^\circ 52.5'. \\
 \therefore c &= 122^\circ 7.5'.
 \end{aligned}$$

Check.

$$\begin{aligned}
 \cos c &= \cos a \cos b. \\
 \log \cos a &= 9.8798 \\
 \log \cos b &= 9.8458 \\
 \log \cos c &= 9.7256
 \end{aligned}$$

$$\begin{aligned}
 7. \quad & \overline{\tan A} = \overline{\frac{\tan a}{\sin b}} \\
 & \quad + \\
 & \log \tan a = 9.5611 \\
 & \log \sin b = 9.7941 \\
 & \log \tan A = 9.7670 \\
 & 180^\circ - A = 30^\circ 19'. \\
 & \therefore A = 149^\circ 41'.
 \end{aligned}$$

$$\begin{aligned}
 & \tan B = \frac{\tan b}{\sin a} \\
 & \log \tan b = 9.9006 \\
 & \log \sin a = 9.5341 \\
 & \log \tan B = 0.3665 \\
 & \therefore B = 66^\circ 43.8'.
 \end{aligned}$$

$$\begin{aligned}
 8. \quad & \sin A = \frac{\cos B}{\cos b} \\
 & \log \cos B = 9.2397 \\
 & \log \cos b = 9.5798 \\
 & \log \sin A = 9.6599 \\
 & \therefore A = 27^\circ 11.6', \\
 & \quad \text{or } 152^\circ 48.4'.
 \end{aligned}$$

$$\begin{aligned}
 & \sin a = \frac{\tan b}{\tan B} \\
 & \log \tan b = 0.3864 \\
 & \log \tan B = 0.7537 \\
 & \log \sin a = 9.6327 \\
 & \therefore a = 25^\circ 25.2', \\
 & \quad \text{or } 154^\circ 34.8'.
 \end{aligned}$$

$$\begin{aligned}
 \text{Ans. } 1. A &= 27^\circ 11.6', a = 25^\circ 25.2', c = 69^\circ 54'. \\
 2. A &= 152^\circ 48.4', a = 154^\circ 34.8', c = 110^\circ 6'.
 \end{aligned}$$

$$\begin{aligned}
 9. \quad & \overline{\tan A} = \overline{\frac{\cot B}{\cos c}} \\
 & \quad + \\
 & \log \cot B = 9.6064 \\
 & \log \cos c = 9.1525 \\
 & \log \tan A = 0.4539 \\
 & 180^\circ - A = 70^\circ 37.5'. \\
 & \therefore A = 109^\circ 22.5'.
 \end{aligned}$$

$$\begin{aligned}
 & \overline{\cos c} = \overline{\cos a \cos b} \\
 & \log \cos a = 9.9730 \\
 & \log \cos b = 9.8935 \\
 & \log \cos c = 9.8665 \\
 & 180^\circ - c = 42^\circ 40'. \\
 & \therefore c = 137^\circ 20'.
 \end{aligned}$$

Check.

$$\begin{aligned}
 \cos c \tan A \tan B &= 1. \\
 \log \cos c &= 9.8665 \\
 \log \tan A &= 9.7670 \\
 \log \tan B &= 0.3665 \\
 \log 1 &= 0.0000
 \end{aligned}$$

$$\begin{aligned}
 \sin c &= \frac{\sin b}{\sin B} \\
 \log \sin b &= 9.9661 \\
 \log \sin B &= 9.9934 \\
 \log \sin c &= 9.9727 \\
 \therefore c &= 69^\circ 54', \\
 & \quad \text{or } 110^\circ 6'.
 \end{aligned}$$

Check.

$$\begin{aligned}
 \sin A &= \frac{\sin a}{\sin c} \\
 \log \sin a &= 9.6327 \\
 \log \sin c &= 9.9727 \\
 \log \sin A &= 9.6600
 \end{aligned}$$

$$\sin b = \sin B \sin c.$$

$$\log \sin B = 9.9672$$

$$\log \sin c = 9.9956$$

$$\log \sin b = 9.9628$$

$$180^\circ - b = 66^\circ 38'.$$

$$\therefore b = 113^\circ 22'.$$

Check.

$$\tan A = \frac{\tan a}{\sin b}.$$

$$\log \tan a = 0.4167$$

$$\log \sin b = 9.9628$$

$$\log \tan A = 0.4539$$

10.

$$\cos A = \cos a \sin B.$$

$$\log \cos a = 9.6856$$

$$\log \sin B = 9.9203$$

$$\log \cos A = 9.6059$$

$$\therefore A = 66^\circ 12.1'.$$

$$\tan b = \sin a \tan B.$$

$$\log \sin a = 9.9418$$

$$\log \tan B = 0.1765$$

$$\log \tan b = 0.1183$$

$$180^\circ - b = 52^\circ 42.6'.$$

$$\therefore b = 127^\circ 17.4'.$$

$$\tan c = \frac{\tan a}{\cos B}.$$

$$\log \tan a = 0.2562$$

$$\log \cos B = 9.7438$$

$$\log \tan c = 0.5124$$

$$180^\circ - c = 72^\circ 54.9'.$$

$$\therefore c = 107^\circ 5.1'.$$

Check.

$$\cos A = \frac{\tan b}{\tan c}.$$

$$\log \tan b = 0.1183$$

$$\log \tan c = 0.5124$$

$$\log \cos A = 9.6059$$

11.

$$\tan A = \frac{\tan a}{\sin b}.$$

$$\log \tan a = 0.2683$$

$$\log \sin b = 9.7675$$

$$\log \tan A = 0.5008$$

$$\therefore A = 72^\circ 28.9'.$$

$$\tan B = \frac{\tan b}{\sin a}.$$

$$\log \tan b = 9.8586$$

$$\log \sin a = 9.9446$$

$$\log \tan B = 9.9140$$

$$180^\circ - B = 39^\circ 21.9'.$$

$$\therefore B = 140^\circ 38.1'.$$

$$\cos c = \cos a \cos b.$$

$$\log \cos a = 9.6763$$

$$\log \cos b = 9.9089$$

$$\log \cos c = 9.5852$$

$$180^\circ - c = 67^\circ 22.3'.$$

$$\therefore c = 112^\circ 37.7'.$$

Check.

$$\cos c \tan A \tan B = 1.$$

$$\log \cos c = 9.5852$$

$$\log \tan A = 0.5008$$

$$\log \tan B = 9.9140$$

$$\log 1 = 0.0000$$

$$\begin{array}{rcl}
 12. & \sin B = \frac{\overset{+}{\cos A}}{\overset{-}{\cos a}} & \\
 & \log \cos A = 9.2324 & \\
 & \log \cos a = 9.5736 & \\
 & \log \sin B = 9.6588 & \\
 & \therefore B = 27^\circ 7.2', & \\
 & \text{or } 152^\circ 52.8'. &
 \end{array}$$

$$\begin{array}{rcl}
 & \sin b = \frac{\overset{+}{\tan a}}{\overset{-}{\tan A}} & \\
 & \log \tan a = 0.3936 & \\
 & \log \tan A = 0.7611 & \\
 & \log \sin b = 9.6325 & \\
 & \therefore b = 25^\circ 24.4', & \\
 & \text{or } 154^\circ 35.6'. &
 \end{array}$$

Ans. 1. $B = 27^\circ 7.2'$, $b = 25^\circ 24.4'$, $c = 109^\circ 46'$.
 2. $B = 152^\circ 52.8'$, $b = 154^\circ 35.6'$, $c = 70^\circ 14'$.

$$\begin{array}{rcl}
 13. & \cos A = \frac{\overset{-}{\tan b}}{\overset{+}{\tan c}} & \\
 & \log \tan b = 9.4281 & \\
 & \log \tan c = 9.7196 & \\
 & \log \cos A = 9.7085 & \\
 & 180^\circ - A = 59^\circ 15.7'. & \\
 & \therefore A = 120^\circ 44.3'. &
 \end{array}$$

$$\begin{array}{rcl}
 & \cos a = \frac{\overset{-}{\cos c}}{\overset{+}{\cos b}} & \\
 & \log \cos c = 9.9473 & \\
 & \log \cos b = 9.9849 & \\
 & \log \cos a = 9.9624 & \\
 & 180^\circ - a = 23^\circ 30'. & \\
 & \therefore a = 156^\circ 30'. &
 \end{array}$$

$$\begin{array}{rcl}
 \sin c = \frac{\sin a}{\sin A} & & \\
 \log \sin a = 9.9672 & & \\
 \log \sin A = 9.9936 & & \\
 \log \sin c = 9.9736 & & \\
 \therefore c = 70^\circ 14', & & \\
 \text{or } 109^\circ 46'. & &
 \end{array}$$

Check.

$$\begin{array}{rcl}
 \sin B = \frac{\sin b}{\sin c} & & \\
 \log \sin b = 9.6325 & & \\
 \log \sin c = 9.9736 & & \\
 \log \sin B = 9.6589 & &
 \end{array}$$

$$\begin{array}{rcl}
 \sin B = \frac{\sin b}{\sin c} & & \\
 \log \sin b = 9.4130 & & \\
 \log \sin c = 9.6668 & & \\
 \log \sin B = 9.7462 & & \\
 \therefore B = 33^\circ 52.6'. & &
 \end{array}$$

Check.

$$\begin{array}{rcl}
 \sin B = \frac{\cos A}{\cos a} & & \\
 \log \cos A = 9.7085 & & \\
 \log \cos a = 9.9624 & & \\
 \log \sin B = 9.7461 & &
 \end{array}$$

$$\begin{aligned}
 14. \quad \cos a &= \frac{\cos A}{\sin B} \\
 \log \cos A &= 9.6573 \\
 \log \sin B &= 9.7801 \\
 \log \cos a &= 9.8772 \\
 \therefore a &= 41^\circ 5.5'.
 \end{aligned}$$

$$\begin{aligned}
 \cos b &= \frac{\cos B}{\sin A} \\
 \log \cos B &= 9.9019 \\
 \log \sin A &= 9.9498 \\
 \log \cos b &= 9.9521 \\
 \therefore b &= 26^\circ 25'.
 \end{aligned}$$

$$\begin{aligned}
 15. \quad \sin a &= \sin A \sin c. \\
 \log \sin A &= 9.9809 \\
 \log \sin c &= 9.9589 \\
 \log \sin a &= 9.9398 \\
 \therefore a &= 60^\circ 31.4'.
 \end{aligned}$$

$$\begin{aligned}
 \tan B &= \frac{\cot A}{\cos c} \\
 \log \cot A &= 9.4822 \\
 \log \cos c &= 9.6183 \\
 \log \tan B &= 9.8639 \\
 180^\circ - B &= 36^\circ 10'. \\
 \therefore B &= 143^\circ 50'.
 \end{aligned}$$

$$\begin{aligned}
 16. \quad \sin A &= \frac{\cos B}{\cos b} \\
 \log \cos B &= 9.9129 \\
 \log \cos b &= 9.9213 \\
 \log \sin A &= 9.9916 \\
 \therefore A &= 78^\circ 46.7', \\
 &\text{or } 101^\circ 13.3'.
 \end{aligned}$$

$$\begin{aligned}
 \cos c &= \cot A \cot B. \\
 \log \cot A &= 9.7075 \\
 \log \cot B &= 0.1219 \\
 \log \cos c &= 9.8294 \\
 \therefore c &= 47^\circ 32.1'.
 \end{aligned}$$

Check.

$$\begin{aligned}
 \cos c &= \cos a \cos b. \\
 \log \cos a &= 9.8772 \\
 \log \cos b &= 9.9521 \\
 \log \cos c &= 9.8293
 \end{aligned}$$

$$\begin{aligned}
 - \quad + \quad - \\
 \tan b &= \cos A \tan c. \\
 \log \cos A &= 9.4630 \\
 \log \tan c &= 0.3406 \\
 \log \tan b &= 9.8036 \\
 180^\circ - b &= 32^\circ 27.9'. \\
 \therefore b &= 147^\circ 32.1'.
 \end{aligned}$$

Check.

$$\begin{aligned}
 \tan B &= \frac{\tan b}{\sin a} \\
 \log \tan b &= 9.8036 \\
 \log \sin a &= 9.9398 \\
 \log \tan B &= 9.8638
 \end{aligned}$$

$$\begin{aligned}
 \sin a &= \frac{\tan b}{\tan B} \\
 \log \tan b &= 9.8202 \\
 \log \tan B &= 9.8468 \\
 \log \sin a &= 9.9734 \\
 \therefore a &= 70^\circ 10', \\
 &\text{or } 109^\circ 50'.
 \end{aligned}$$

$$\sin c = \frac{\sin b}{\sin B}.$$

$$\log \sin b = 9.7415$$

$$\log \sin B = \underline{9.7597}$$

$$\log \sin c = \underline{9.9818}$$

$$\therefore c = 73^\circ 32.5',$$

$$\text{or } 106^\circ 27.5'.$$

$$\text{Ans. } 1. A = 78^\circ 46.7', a = 70^\circ 10', c = 106^\circ 27.5'.$$

$$2. A = 101^\circ 13.3', a = 109^\circ 50', c = 73^\circ 32.5'.$$

Check.

$$\sin A = \frac{\sin a}{\sin c}.$$

$$\log \sin a = 9.9734$$

$$\log \sin c = \underline{9.9818}$$

$$\log \sin A = \underline{9.9916}$$

$$17. \quad \tan A = \frac{\cot B}{\cos c}.$$

$$\log \cot B = 9.5998$$

$$\log \cos c = \underline{9.8291}$$

$$\log \tan A = \underline{9.7707}$$

$$\therefore A = 30^\circ 32.1'.$$

$$\sin b = \sin B \sin c.$$

$$\log \sin B = 9.9681$$

$$\log \sin c = \underline{9.8681}$$

$$\log \sin b = \underline{9.8362}$$

$$\therefore b = 43^\circ 17.9'.$$

Check.

$$\tan A = \frac{\tan a}{\sin b}.$$

$$\log \tan a = 9.6068$$

$$\log \sin b = \underline{9.8362}$$

$$\log \tan A = \underline{9.7706}$$

$$\tan a = \cos B \tan c.$$

$$\log \cos B = 9.5679$$

$$\log \tan c = \underline{0.0389}$$

$$\log \tan a = \underline{9.6068}$$

$$\therefore a = 22^\circ 1.1'.$$

$$18. \quad \tan a = \tan A \sin b.$$

$$\log \tan A = 9.5152$$

$$\log \sin b = \underline{9.8769}$$

$$\log \tan a = \underline{9.3921}$$

$$180^\circ - a = 13^\circ 51.3'.$$

$$\therefore a = 166^\circ 8.7'.$$

$$\tan c = \frac{\tan b}{\cos A}.$$

$$\log \tan b = 0.0588$$

$$\log \cos A = \underline{9.9779}$$

$$\log \tan c = \underline{0.0809}$$

$$\therefore c = 50^\circ 18.4'.$$

$$\cos B = \sin A \cos b.$$

$$\log \sin A = 9.4931$$

$$\log \cos b = \underline{9.8181}$$

$$\log \cos B = \underline{9.3112}$$

$$180^\circ - B = 78^\circ 11.1'.$$

$$\therefore B = 101^\circ 48.9'.$$

Check.

$$\cos B = \frac{\tan a}{\tan c}.$$

$$\log \tan a = 9.3921$$

$$\log \tan c = \underline{0.0809}$$

$$\log \cos B = \underline{9.3112}$$

$$\begin{array}{rcl}
 19. & \overline{\tan A} = \frac{\overline{\tan a}}{\overline{\sin b}} & \overline{\tan B} = \frac{\overline{\tan b}}{\overline{\sin a}} \\
 & + & + \\
 & \log \tan a = 0.3634 & \log \tan b = 0.4207 \\
 & \log \sin b = 9.9707 & \log \sin a = 9.9626 \\
 & \hline
 & \log \tan A = 0.3927 & \log \tan B = 0.4581 \\
 & 180^\circ - A = 67^\circ 57.5'. & 180^\circ - B = 70^\circ 48'. \\
 & \therefore A = 112^\circ 2.5'. & \therefore B = 109^\circ 12'.
 \end{array}$$

$$\begin{array}{rcl}
 & + & - & - \\
 & \cos c = \cos a \cos b. & & \text{Check.} \\
 & \log \cos a = 9.5993 & & \cos c \tan A \tan B = 1. \\
 & \log \cos b = 9.5500 & & \log \cos c = 9.1493 \\
 & \hline
 & \log \cos c = 9.1493 & & \log \tan A = 0.3927 \\
 & \therefore c = 81^\circ 53.6'. & & \log \tan B = 0.4581 \\
 & & & \hline
 & & & \log 1 = 0.0001
 \end{array}$$

$$\begin{array}{rcl}
 20. & \overline{\cos A} = \overline{\cos a} \overline{\sin B} & \overline{\tan c} = \frac{\overline{\tan a}}{\overline{\cos B}} \\
 & \log \cos a = 9.8652 & + \\
 & \log \sin B = 9.9846 & \log \tan a = 9.9674 \\
 & \hline
 & \log \cos A = 9.8498 & \log \cos B = 9.4172 \\
 & 180^\circ - A = 44^\circ 57.5'. & \hline
 & \therefore A = 135^\circ 2.5'. & \log \tan c = 0.5502 \\
 & & 180^\circ - c = 74^\circ 16'. \\
 & & \therefore c = 105^\circ 44'.
 \end{array}$$

$$\begin{array}{rcl}
 & \tan b = \sin a \tan B. & \text{Check.} \\
 & \log \sin a = 9.8325 & \cos A = \frac{\tan b}{\tan c} \\
 & \log \tan B = 0.5674 & \log \tan b = 0.3999 \\
 & \hline
 & \log \tan b = 0.3999 & \log \tan c = 0.5502 \\
 & \therefore b = 68^\circ 17.3'. & \hline
 & & \log \cos A = 9.8497
 \end{array}$$

$$\begin{array}{rcl}
 21. & \overline{\cos a} = \frac{\overline{\cos A}}{\overline{\sin B}} & \overline{\cos b} = \frac{\overline{\cos B}}{\overline{\sin A}} \\
 & + & + \\
 & \log \cos A = 9.9129 & \log \cos B = 9.2896 \\
 & \log \sin B = 9.9916 & \log \sin A = 9.7597 \\
 & \hline
 & \log \cos a = 9.9213 & \log \cos b = 9.5299 \\
 & 180^\circ - a = 33^\circ 27.5'. & 180^\circ - b = 70^\circ 11.9'. \\
 & \therefore a = 146^\circ 32.5'. & \therefore b = 109^\circ 48.1'.
 \end{array}$$

$$\begin{array}{c} + \quad - \quad - \\ \cos c = \cot A \cot B. \end{array}$$

$$\log \cot A = 0.1532$$

$$\log \cot B = \underline{9.2980}$$

$$\log \cos c = 9.4512$$

$$\therefore c = 73^\circ 35'.$$

Check.

$$\cos c = \cos a \cos b.$$

$$\log \cos a = 9.9213$$

$$\log \cos b = \underline{9.5299}$$

$$\log \cos c = 9.4512$$

22.

$$\sin A = \frac{\sin a}{\sin c}$$

$$\log \sin a = 9.9710$$

$$\log \sin c = \underline{9.9980}$$

$$\log \sin A = 9.9730$$

$$\therefore A = 70^\circ.$$

$$\cos b = \frac{\cos c}{\cos a}$$

$$\log \cos c = 8.9855$$

$$\log \cos a = \underline{9.5484}$$

$$\log \cos b = 9.4371$$

$$\therefore b = 74^\circ 7.3'.$$

$$\cos B = \frac{\tan a}{\tan c}$$

$$\log \tan a = 0.4226$$

$$\log \tan c = \underline{1.0125}$$

$$\log \cos B = 9.4101$$

$$\therefore B = 75^\circ 6.2'.$$

Check.

$$\sin A = \frac{\cos B}{\cos b}$$

$$\log \cos B = 9.4101$$

$$\log \cos b = \underline{9.4371}$$

$$\log \sin A = 9.9730$$

Art. 154.—Page 113.

2. In the polar triangle, $a' = 41^\circ$ and $B' = 37^\circ$.

$$\cos A' = \cos a' \sin B'.$$

$$\log \cos a' = 9.8778$$

$$\log \sin B' = \underline{9.7795}$$

$$\log \cos A' = 9.6573$$

$$\therefore A' = 62^\circ 58.8'.$$

$$\tan b' = \sin a' \tan B'.$$

$$\log \sin a' = 9.8169$$

$$\log \tan B' = \underline{9.8771}$$

$$\log \tan b' = 9.6940$$

$$\therefore b' = 26^\circ 18.1'.$$

$$\tan c' = \frac{\tan a'}{\cos B'}$$

$$\log \tan a' = 9.9392$$

$$\log \cos B' = \underline{9.9023}$$

$$\log \tan c' = 0.0369$$

$$\therefore c' = 47^\circ 26'.$$

Check.

$$\cos A' = \frac{\tan b'}{\tan c'}$$

$$\log \tan b' = 9.6940$$

$$\log \tan c' = \underline{0.0369}$$

$$\log \cos A' = 9.6571$$

\therefore in the quadrantal triangle,

$$a = 117^\circ 1.2', B = 153^\circ 41.9', \text{ and } C = 132^\circ 34'.$$

3. In the polar triangle, $a' = 134^\circ 30'$ and $b' = 40^\circ 40'$.

$$\begin{array}{rcl}
 \tan A' & = & \frac{\tan a'}{\sin b'} \\
 \log \tan a' & = & 0.0076 \\
 \log \sin b' & = & 9.8140 \\
 \log \tan A' & = & 0.1936 \\
 \therefore 180^\circ - A' & = & 57^\circ 22.1'.
 \end{array}
 \qquad
 \begin{array}{rcl}
 \cos c' & = & \cos a' \cos b' \\
 \log \cos a' & = & 9.8457 \\
 \log \cos b' & = & 9.8800 \\
 \log \cos c' & = & 9.7257 \\
 \therefore 180^\circ - c' & = & 57^\circ 52.5'.
 \end{array}$$

Check.

$$\begin{array}{rcl}
 \tan B' & = & \frac{\tan b'}{\sin a'} \\
 \log \tan b' & = & 9.9341 \\
 \log \sin a' & = & 9.8532 \\
 \log \tan B' & = & 0.0809 \\
 \therefore B' & = & 50^\circ 18.4'.
 \end{array}
 \qquad
 \begin{array}{rcl}
 \cos c' \tan A' \tan B' & = & 1. \\
 \log \cos c' & = & 9.7257 \\
 \log \tan A' & = & 0.1936 \\
 \log \tan B' & = & 0.0809 \\
 \log 1 & = & 0.0002
 \end{array}$$

\therefore in the quadrantal triangle,

$$a = 57^\circ 22.1', \quad b = 129^\circ 41.6', \quad \text{and} \quad C = 57^\circ 52.5'.$$

4. In the polar triangle, $A' = 149^\circ 40'$ and $c' = 137^\circ 20'$.

$$\begin{array}{rcl}
 \sin a' & = & \sin A' \sin c' \\
 \log \sin A' & = & 9.7033 \\
 \log \sin c' & = & 9.8311 \\
 \log \sin a' & = & 9.5344 \\
 \therefore 180^\circ - a' & = & 20^\circ 0.9'.
 \end{array}
 \qquad
 \begin{array}{rcl}
 \tan b' & = & \cos A' \tan c' \\
 \log \cos A' & = & 9.9361 \\
 \log \tan c' & = & 9.9646 \\
 \log \tan b' & = & 9.9007 \\
 \therefore b' & = & 38^\circ 30.4'.
 \end{array}$$

Check.

$$\begin{array}{rcl}
 \tan B' & = & \frac{\cot A'}{\cos c'} \\
 \log \cot A' & = & 0.2327 \\
 \log \cos c' & = & 9.8665 \\
 \log \tan B' & = & 0.3662 \\
 \therefore B' & = & 66^\circ 42.9'.
 \end{array}
 \qquad
 \begin{array}{rcl}
 \tan B' & = & \frac{\tan b'}{\sin a'} \\
 \log \tan b' & = & 9.9007 \\
 \log \sin a' & = & 9.5344 \\
 \log \tan B' & = & 0.3663
 \end{array}$$

\therefore in the quadrantal triangle,

$$A = 20^\circ 0.9', \quad b = 113^\circ 17.1', \quad \text{and} \quad B = 141^\circ 29.6'.$$

5. In the polar triangle, $b' = 109^\circ 48'$ and $c' = 73^\circ 35'$.

$$\begin{array}{rcl} \cos a' & = & \frac{\cos c'}{\cos b'} \\ \log \cos c' & = & 9.4512 \\ \log \cos b' & = & 9.5299 \\ \log \cos a' & = & 9.9213 \\ \therefore 180^\circ - a' & = & 33^\circ 27.5'. \end{array} \qquad \begin{array}{rcl} \cos A' & = & \frac{\tan b'}{\tan c'} \\ \log \tan b' & = & 0.4437 \\ \log \tan c' & = & 0.5307 \\ \log \cos A' & = & 9.9130 \\ \therefore 180^\circ - A' & = & 35^\circ 4.4'. \end{array}$$

$$\begin{array}{rcl} \sin B' & = & \frac{\sin b'}{\sin c'} \\ \log \sin b' & = & 9.9735 \\ \log \sin c' & = & 9.9819 \\ \log \sin B' & = & 9.9916 \\ \therefore 180^\circ - B' & = & 78^\circ 46.7'. \end{array} \qquad \begin{array}{l} \text{Check.} \\ \sin B' = \frac{\cos A'}{\cos a'} \\ \log \cos A' = 9.9130 \\ \log \cos a = 9.9213 \\ \log \sin B' = 9.9917 \end{array}$$

\therefore in the quadrantal triangle,

$$a = 35^\circ 4.4', \quad A = 33^\circ 27.5', \quad \text{and} \quad b = 78^\circ 46.7'.$$

6. In the polar triangle, $a' = 74^\circ 7'$ and $A' = 75^\circ 6'$.

$$\begin{array}{rcl} \sin b' & = & \frac{\tan a'}{\tan A'} \\ \log \tan a' & = & 0.5459 \\ \log \tan A' & = & 0.5750 \\ \log \sin b' & = & 9.9709 \\ \therefore b' & = & 69^\circ 16', \\ & \text{or} & 110^\circ 44'. \end{array} \qquad \begin{array}{rcl} \sin B' & = & \frac{\cos A'}{\cos a'} \\ \log \cos A' & = & 9.4102 \\ \log \cos a' & = & 9.4372 \\ \log \sin B' & = & 9.9730 \\ \therefore B' & = & 70^\circ, \\ & \text{or} & 110^\circ. \end{array}$$

$$\begin{array}{rcl} \sin c' & = & \frac{\sin a'}{\sin A'} \\ \log \sin a' & = & 9.9831 \\ \log \sin A' & = & 9.9851 \\ \log \sin c' & = & 9.9980 \\ \therefore c' & = & 84^\circ 30', \\ & \text{or} & 95^\circ 30'. \end{array} \qquad \begin{array}{l} \text{Check.} \\ \sin B' = \frac{\sin b'}{\sin c'} \\ \log \sin b' = 9.9709 \\ \log \sin c' = 9.9980 \\ \log \sin B' = 9.9729 \end{array}$$

\therefore in the quadrantal triangle,

1. $B = 69^\circ 16'$, $b = 70^\circ$, and $C = 84^\circ 30'$.
2. $B = 110^\circ 44'$, $b = 110^\circ$, and $C = 95^\circ 30'$.

CHAPTER XII.

Art. 167. — Page 125.

2. Here $\frac{1}{2}(A - B) = 18^\circ 30'$, $\frac{1}{2}(A + B) = 59^\circ 30'$, $\frac{1}{2}c = 54^\circ$.

$$\tan \frac{1}{2}(a - b) = \frac{\sin \frac{1}{2}(A - B)}{\sin \frac{1}{2}(A + B)} \tan \frac{1}{2}c. \quad \tan \frac{1}{2}(a + b) = \frac{\cos \frac{1}{2}(A - B)}{\cos \frac{1}{2}(A + B)} \tan \frac{1}{2}c.$$

$$\log \sin \frac{1}{2}(A - B) = 9.5015$$

$$\log \cos \frac{1}{2}(A - B) = 9.9770$$

$$\log \csc \frac{1}{2}(A + B) = 0.0647$$

$$\log \sec \frac{1}{2}(A + B) = 0.2945$$

$$\log \tan \frac{1}{2}c = 0.1387$$

$$\log \tan \frac{1}{2}c = 0.1387$$

$$\log \tan \frac{1}{2}(a - b) = 9.7049$$

$$\log \tan \frac{1}{2}(a + b) = 0.4102$$

$$\therefore \frac{1}{2}(a - b) = 26^\circ 52.8'.$$

$$\therefore \frac{1}{2}(a + b) = 68^\circ 45'.$$

$$\therefore a = \frac{1}{2}(a + b) + \frac{1}{2}(a - b) = 95^\circ 37.8',$$

and .

$$b = \frac{1}{2}(a + b) - \frac{1}{2}(a - b) = 41^\circ 52.2'.$$

$$\cot \frac{1}{2}C = \frac{\sin \frac{1}{2}(a + b)}{\sin \frac{1}{2}(a - b)} \tan \frac{1}{2}(A - B).$$

$$\log \sin \frac{1}{2}(a + b) = 9.9694$$

$$\log \csc \frac{1}{2}(a - b) = 0.3447$$

$$\log \tan \frac{1}{2}(A - B) = 9.5245$$

$$\log \cot \frac{1}{2}C = 9.8386$$

$$\frac{1}{2}C = 55^\circ 24.4'$$

$$\therefore C = 110^\circ 48.8'.$$

3. Here $\frac{1}{2}(B - C) = 42^\circ 30'$, $\frac{1}{2}(B + C) = 92^\circ 30'$, $\frac{1}{2}a = 35^\circ 10'$.

$$\tan \frac{1}{2}(b - c) = \frac{\sin \frac{1}{2}(B - C)}{\sin \frac{1}{2}(B + C)} \tan \frac{1}{2}a. \quad \tan \frac{1}{2}(b + c) = \frac{\cos \frac{1}{2}(B - C)}{\cos \frac{1}{2}(B + C)} \tan \frac{1}{2}a.$$

$$\log \sin \frac{1}{2}(B - C) = 9.8297$$

$$\log \cos \frac{1}{2}(B - C) = 9.8676$$

$$\log \csc \frac{1}{2}(B + C) = 0.0004$$

$$\log \sec \frac{1}{2}(B + C) = 1.3603$$

$$\log \tan \frac{1}{2}a = 9.8479$$

$$\log \tan \frac{1}{2}a = 9.8479$$

$$\log \tan \frac{1}{2}(b - c) = 9.6780$$

$$\log \tan \frac{1}{2}(b + c) = 1.0758$$

$$\therefore \frac{1}{2}(b - c) = 25^\circ 28.5'.$$

$$180^\circ - \frac{1}{2}(b + c) = 85^\circ 11.9'$$

$$\therefore \frac{1}{2}(b + c) = 94^\circ 48.1'.$$

$$\therefore b = \frac{1}{2}(b + c) + \frac{1}{2}(b - c) = 120^\circ 16.6',$$

and

$$c = \frac{1}{2}(b + c) - \frac{1}{2}(b - c) = 69^\circ 19.6'.$$

$$\cot \frac{1}{2} A = \frac{\sin \frac{1}{2} (b+c)}{\sin \frac{1}{2} (b-c)} \tan \frac{1}{2} (B-C).$$

$$\log \sin \frac{1}{2} (b+c) = 9.9985$$

$$\log \csc \frac{1}{2} (b-c) = 0.3664$$

$$\log \tan \frac{1}{2} (B-C) = \underline{9.9621}$$

$$\log \cot \frac{1}{2} A = \underline{0.3270}$$

$$\frac{1}{2} A = 25^\circ 13.1'$$

$$\therefore A = 50^\circ 26.2'.$$

4. Here $\frac{1}{2} (C-A) = 45^\circ 20'$, $\frac{1}{2} (C+A) = 77^\circ$, $\frac{1}{2} b = 20^\circ 20'$.

$$\tan \frac{1}{2} (c-a) = \frac{\sin \frac{1}{2} (C-A)}{\sin \frac{1}{2} (C+A)} \tan \frac{1}{2} b. \quad \tan \frac{1}{2} (c+a) = \frac{\cos \frac{1}{2} (C-A)}{\cos \frac{1}{2} (C+A)} \tan \frac{1}{2} b.$$

$$\log \sin \frac{1}{2} (C-A) = 9.8520$$

$$\log \cos \frac{1}{2} (C-A) = 9.8469$$

$$\log \csc \frac{1}{2} (C+A) = 0.0113$$

$$\log \sec \frac{1}{2} (C+A) = 0.6479$$

$$\log \tan \frac{1}{2} b = \underline{9.5689}$$

$$\log \tan \frac{1}{2} b = \underline{9.5689}$$

$$\log \tan \frac{1}{2} (c-a) = \underline{9.4322}$$

$$\log \tan \frac{1}{2} (c+a) = \underline{0.0637}$$

$$\therefore \frac{1}{2} (c-a) = 15^\circ 8.2'.$$

$$\therefore \frac{1}{2} (c+a) = 49^\circ 11.2'.$$

$$\therefore a = \frac{1}{2} (c+a) - \frac{1}{2} (c-a) = 34^\circ 3',$$

and

$$c = \frac{1}{2} (c+a) + \frac{1}{2} (c-a) = 64^\circ 19.4'.$$

$$\cot \frac{1}{2} B = \frac{\sin \frac{1}{2} (c+a)}{\sin \frac{1}{2} (c-a)} \tan \frac{1}{2} (C-A).$$

$$\log \sin \frac{1}{2} (c+a) = 9.8790$$

$$\log \csc \frac{1}{2} (c-a) = 0.5831$$

$$\log \tan \frac{1}{2} (C-A) = \underline{0.0051}$$

$$\log \cot \frac{1}{2} B = \underline{0.4672}$$

$$\frac{1}{2} B = 18^\circ 49.8'$$

$$\therefore B = 37^\circ 39.6'.$$

5. Here $\frac{1}{2} (B-A) = 18^\circ 47'$, $\frac{1}{2} (B+A) = 126^\circ 59'$, $\frac{1}{2} c = 63^\circ 16'$.

$$\tan \frac{1}{2} (b-a) = \frac{\sin \frac{1}{2} (B-A)}{\sin \frac{1}{2} (B+A)} \tan \frac{1}{2} c. \quad \tan \frac{1}{2} (b+a) = \frac{\cos \frac{1}{2} (B-A)}{\cos \frac{1}{2} (B+A)} \tan \frac{1}{2} c.$$

$$\log \sin \frac{1}{2} (B-A) = 9.5079$$

$$\log \cos \frac{1}{2} (B-A) = 9.9762$$

$$\log \csc \frac{1}{2} (B+A) = 0.0976$$

$$\log \sec \frac{1}{2} (B+A) = 0.2207$$

$$\log \tan \frac{1}{2} c = \underline{0.2979}$$

$$\log \tan \frac{1}{2} c = \underline{0.2979}$$

$$\log \tan \frac{1}{2} (b-a) = \underline{9.9034}$$

$$\log \tan \frac{1}{2} (b+a) = \underline{0.4948}$$

$$\therefore \frac{1}{2} (b-a) = 38^\circ 40.8'.$$

$$180^\circ - \frac{1}{2} (b+a) = 72^\circ 15.2'$$

$$\therefore \frac{1}{2} (b+a) = 107^\circ 44.8'.$$

$$\therefore a = \frac{1}{2} (b+a) - \frac{1}{2} (b-a) = 69^\circ 4',$$

and

$$b = \frac{1}{2} (b+a) + \frac{1}{2} (b-a) = 146^\circ 25.6'.$$

$$\cot \frac{1}{2} C = \frac{\sin \frac{1}{2} (b + a)}{\sin \frac{1}{2} (b - a)} \tan \frac{1}{2} (B - A).$$

$$\log \sin \frac{1}{2} (b + a) = 9.9788$$

$$\log \csc \frac{1}{2} (b - a) = 0.2042$$

$$\log \tan \frac{1}{2} (B - A) = 9.5316$$

$$\log \cot \frac{1}{2} C = 9.7146$$

$$\frac{1}{2} C = 62^\circ 36.1'$$

$$\therefore C = 125^\circ 12.2'.$$

Art. 168.—Page 126.

$$2. \text{ Here } \frac{1}{2} (a - b) = 12^\circ 30', \frac{1}{2} (a + b) = 59^\circ 30', \frac{1}{2} C = 16^\circ 30'.$$

$$\tan \frac{1}{2} (A - B) = \frac{\sin \frac{1}{2} (a - b)}{\sin \frac{1}{2} (a + b)} \cot \frac{1}{2} C. \quad \tan \frac{1}{2} (A + B) = \frac{\cos \frac{1}{2} (a - b)}{\cos \frac{1}{2} (a + b)} \cot \frac{1}{2} C.$$

$$\log \sin \frac{1}{2} (a - b) = 9.3353$$

$$\log \cos \frac{1}{2} (a - b) = 9.9896$$

$$\log \csc \frac{1}{2} (a + b) = 0.0647$$

$$\log \sec \frac{1}{2} (a + b) = 0.2945$$

$$\log \cot \frac{1}{2} C = 0.5284$$

$$\log \cot \frac{1}{2} C = 0.5284$$

$$\log \tan \frac{1}{2} (A - B) = 9.9284$$

$$\log \tan \frac{1}{2} (A + B) = 0.8125$$

$$\therefore \frac{1}{2} (A - B) = 40^\circ 18'.$$

$$\therefore \frac{1}{2} (A + B) = 81^\circ 14.8'.$$

$$\therefore A = \frac{1}{2} (A + B) + \frac{1}{2} (A - B) = 121^\circ 32.8',$$

and

$$B = \frac{1}{2} (A + B) - \frac{1}{2} (A - B) = 40^\circ 56.8'.$$

$$\tan \frac{1}{2} c = \frac{\sin \frac{1}{2} (A + B)}{\sin \frac{1}{2} (A - B)} \tan \frac{1}{2} (a - b).$$

$$\log \sin \frac{1}{2} (A + B) = 9.9949$$

$$\log \csc \frac{1}{2} (A - B) = 0.1892$$

$$\log \tan \frac{1}{2} (a - b) = 9.3458$$

$$\log \tan \frac{1}{2} c = 9.5299$$

$$\frac{1}{2} c = 18^\circ 42.9'$$

$$\therefore c = 37^\circ 25.8'.$$

$$3. \text{ Here } \frac{1}{2} (a - c) = 19^\circ, \frac{1}{2} (a + c) = 79^\circ, \frac{1}{2} B = 55^\circ.$$

$$\tan \frac{1}{2} (A - C) = \frac{\sin \frac{1}{2} (a - c)}{\sin \frac{1}{2} (a + c)} \cot \frac{1}{2} B. \quad \tan \frac{1}{2} (A + C) = \frac{\cos \frac{1}{2} (a - c)}{\cos \frac{1}{2} (a + c)} \cot \frac{1}{2} B.$$

$$\log \sin \frac{1}{2} (a - c) = 9.5126$$

$$\log \cos \frac{1}{2} (a - c) = 9.9757$$

$$\log \csc \frac{1}{2} (a + c) = 0.0081$$

$$\log \sec \frac{1}{2} (a + c) = 0.7194$$

$$\log \cot \frac{1}{2} B = 9.8452$$

$$\log \cot \frac{1}{2} B = 9.8452$$

$$\log \tan \frac{1}{2} (A - C) = 9.3659$$

$$\log \tan \frac{1}{2} (A + C) = 0.5403$$

$$\therefore \frac{1}{2} (A - C) = 13^\circ 4.4'.$$

$$\therefore \frac{1}{2} (A + C) = 73^\circ 55.3'.$$

$$\therefore A = \frac{1}{2} (A + C) + \frac{1}{2} (A - C) = 86^\circ 59.7',$$

and

$$C = \frac{1}{2} (A + C) - \frac{1}{2} (A - C) = 60^\circ 50.9'.$$

$$\tan \frac{1}{2} b = \frac{\sin \frac{1}{2} (A + C)}{\sin \frac{1}{2} (A - C)} \tan \frac{1}{2} (a - c).$$

$$\log \sin \frac{1}{2} (A + C) = 9.9827$$

$$\log \csc \frac{1}{2} (A - C) = 0.6455$$

$$\log \tan \frac{1}{2} (a - c) = 9.5370$$

$$\log \tan \frac{1}{2} b = 0.1652$$

$$\frac{1}{2} b = 55^\circ 38.5'$$

$$\therefore b = 111^\circ 17'.$$

$$4. \text{ Here } \frac{1}{2} (b - c) = 24^\circ 50', \frac{1}{2} (b + c) = 95^\circ 30', \frac{1}{2} A = 25^\circ.$$

$$\tan \frac{1}{2} (B - C) = \frac{\sin \frac{1}{2} (b - c)}{\sin \frac{1}{2} (b + c)} \cot \frac{1}{2} A.$$

$$\log \sin \frac{1}{2} (b - c) = 9.6232$$

$$\log \csc \frac{1}{2} (b + c) = 0.0020$$

$$\log \cot \frac{1}{2} A = 0.3313$$

$$\log \tan \frac{1}{2} (B - C) = 9.9565$$

$$\therefore \frac{1}{2} (B - C) = 42^\circ 8.1'.$$

$$\tan \frac{1}{2} (B + C) = \frac{\cos \frac{1}{2} (b - c)}{\cos \frac{1}{2} (b + c)} \cot \frac{1}{2} A.$$

$$\log \cos \frac{1}{2} (b - c) = 9.9579$$

$$\log \sec \frac{1}{2} (b + c) = 1.0184$$

$$\log \cot \frac{1}{2} A = 0.3313$$

$$\log \tan \frac{1}{2} (B + C) = 1.3076$$

$$180^\circ - \frac{1}{2} (B + C) = 87^\circ 10.8'$$

$$\therefore \frac{1}{2} (B + C) = 92^\circ 49.2'.$$

$$\therefore B = \frac{1}{2} (B + C) + \frac{1}{2} (B - C) = 134^\circ 57.3',$$

and

$$C = \frac{1}{2} (B + C) - \frac{1}{2} (B - C) = 50^\circ 41.1'.$$

$$\tan \frac{1}{2} a = \frac{\sin \frac{1}{2} (B + C)}{\sin \frac{1}{2} (B - C)} \tan \frac{1}{2} (b - c).$$

$$\log \sin \frac{1}{2} (B + C) = 9.9995$$

$$\log \csc \frac{1}{2} (B - C) = 0.1734$$

$$\log \tan \frac{1}{2} (b - c) = 9.6664$$

$$\log \tan \frac{1}{2} a = 9.8393$$

$$\frac{1}{2} a = 34^\circ 34.4'$$

$$\therefore a = 69^\circ 8.8'.$$

$$5. \text{ Here } \frac{1}{2} (b - a) = 14^\circ 20', \frac{1}{2} (b + a) = 139^\circ 30', \frac{1}{2} C = 70^\circ 10'.$$

$$\tan \frac{1}{2} (B - A) = \frac{\sin \frac{1}{2} (b - a)}{\sin \frac{1}{2} (b + a)} \cot \frac{1}{2} C.$$

$$\log \sin \frac{1}{2} (b - a) = 9.3937$$

$$\log \csc \frac{1}{2} (b + a) = 0.1875$$

$$\log \cot \frac{1}{2} C = 9.5571$$

$$\log \tan \frac{1}{2} (B - A) = 9.1383$$

$$\therefore \frac{1}{2} (B - A) = 7^\circ 49.8'.$$

$$\tan \frac{1}{2} (B + A) = \frac{\cos \frac{1}{2} (b - a)}{\cos \frac{1}{2} (b + a)} \cot \frac{1}{2} C.$$

$$\log \cos \frac{1}{2} (b - a) = 9.9863$$

$$\log \sec \frac{1}{2} (b + a) = 0.1190$$

$$\log \cot \frac{1}{2} C = 9.5571$$

$$\log \tan \frac{1}{2} (B + A) = 9.6624$$

$$180^\circ - \frac{1}{2} (B + A) = 24^\circ 41.2'$$

$$\therefore \frac{1}{2} (B + A) = 155^\circ 18.8'.$$

$$\therefore A = \frac{1}{2} (B + A) - \frac{1}{2} (B - A) = 147^\circ 29',$$

and

$$B = \frac{1}{2} (B + A) + \frac{1}{2} (B - A) = 163^\circ 8.6'.$$

$$\tan \frac{1}{2} c = \frac{\sin \frac{1}{2} (B + A)}{\sin \frac{1}{2} (B - A)} \tan \frac{1}{2} (b - a).$$

$$\log \sin \frac{1}{2} (B + A) = 9.6208$$

$$\log \csc \frac{1}{2} (B - A) = 0.8657$$

$$\log \tan \frac{1}{2} (b - a) = 9.4074$$

$$\log \tan \frac{1}{2} c = 9.8939$$

$$\frac{1}{2} c = 38^\circ 4.2'$$

$$\therefore c = 76^\circ 8.4'.$$

Art. 169.—Page 127.

2. Here $s = 65^\circ 30'$, $s - a = 27^\circ 30'$, $s - b = 14^\circ 30'$, $s - c = 23^\circ 30'$.

$$\log \sin (s - b) = 9.3986$$

$$\log \sin (s - c) = 9.6007$$

$$\log \sin (s - c) = 9.6007$$

$$\log \sin (s - a) = 9.6644$$

$$\log \csc s = 0.0410$$

$$\log \csc s = 0.0410$$

$$\log \csc (s - a) = 0.3356$$

$$\log \csc (s - b) = 0.6014$$

$$\begin{array}{r} 2) 9.3759 \\ \log \tan \frac{1}{2} A = 9.6879 \end{array}$$

$$\begin{array}{r} 2) 9.9075 \\ \log \tan \frac{1}{2} B = 9.9537 \end{array}$$

$$\frac{1}{2} A = 25^\circ 59.1'$$

$$\frac{1}{2} B = 41^\circ 57.2'$$

$$\therefore A = 51^\circ 58.2'.$$

$$\therefore B = 83^\circ 54.4'.$$

$$\log \sin (s - a) = 9.6644$$

$$\log \sin (s - b) = 9.3986$$

$$\log \csc s = 0.0410$$

$$\log \csc (s - c) = 0.3993$$

$$\begin{array}{r} 2) 9.5033 \\ \log \tan \frac{1}{2} C = 9.7516 \end{array}$$

$$\frac{1}{2} C = 29^\circ 26.6'$$

$$\therefore C = 58^\circ 53.2'.$$

3. Here $s = 105^\circ$, $s - a = 4^\circ$, $s - b = 56^\circ$, $s - c = 45^\circ$.

$$\log \sin (s - b) = 9.9186$$

$$\log \sin (s - c) = 9.8495$$

$$\log \sin (s - c) = 9.8495$$

$$\log \sin (s - a) = 8.8436$$

$$\log \csc s = 0.0151$$

$$\log \csc s = 0.0151$$

$$\log \csc (s - a) = 1.1564$$

$$\log \csc (s - b) = 0.0814$$

$$\begin{array}{r} 2) 0.9396 \\ \log \tan \frac{1}{2} A = 0.4698 \end{array}$$

$$\begin{array}{r} 2) 8.7896 \\ \log \tan \frac{1}{2} B = 9.3948 \end{array}$$

$$\frac{1}{2} A = 71^\circ 16.4'$$

$$\frac{1}{2} B = 13^\circ 56.3'$$

$$\therefore A = 142^\circ 32.8'.$$

$$\therefore B = 27^\circ 52.6'.$$

$$\begin{aligned}
 \log \sin (s-a) &= 8.8436 \\
 \log \sin (s-b) &= 9.9186 \\
 \log \csc s &= 0.0151 \\
 \log \csc (s-c) &= 0.1505 \\
 &\quad \underline{2) 8.9278} \\
 \log \tan \frac{1}{2} C &= 9.4639 \\
 \frac{1}{2} C &= 16^\circ 13.6' \\
 \therefore C &= 32^\circ 27.2'.
 \end{aligned}$$

4. Here $s = 96^\circ$, $s - a = 35^\circ$, $s - b = 57^\circ$, $s - c = 4^\circ$.

$$\begin{array}{ll}
 \log \sin (s-b) = 9.9236 & \log \sin (s-c) = 8.8436 \\
 \log \sin (s-c) = 8.8436 & \log \sin (s-a) = 9.7586 \\
 \log \csc s = 0.0024 & \log \csc s = 0.0024 \\
 \log \csc (s-a) = 0.2414 & \log \csc (s-b) = 0.0764 \\
 & \quad \underline{2) 8.6810} \\
 \log \tan \frac{1}{2} A = 9.5055 & \log \tan \frac{1}{2} B = 9.3405 \\
 \frac{1}{2} A = 17^\circ 45.6' & \frac{1}{2} B = 12^\circ 21.3' \\
 \therefore A = 35^\circ 31.2' & \therefore B = 24^\circ 42.6'.
 \end{array}$$

$$\begin{aligned}
 \log \sin (s-a) &= 9.7586 \\
 \log \sin (s-b) &= 9.9236 \\
 \log \csc s &= 0.0024 \\
 \log \csc (s-c) &= 1.1564 \\
 &\quad \underline{2) 0.8410} \\
 \log \tan \frac{1}{2} C &= 0.4205 \\
 \frac{1}{2} C &= 69^\circ 12.4' \\
 \therefore C &= 138^\circ 24.8'.
 \end{aligned}$$

5. Here $s = 107^\circ 10'$, $s - a = 44^\circ 50'$, $s - b = 53^\circ$, $s - c = 9^\circ 20'$.

$$\begin{array}{ll}
 \log \sin (s-b) = 9.9023 & \log \sin (s-c) = 9.2100 \\
 \log \sin (s-c) = 9.2100 & \log \sin (s-a) = 9.8482 \\
 \log \csc s = 0.0198 & \log \csc s = 0.0198 \\
 \log \csc (s-a) = 0.1518 & \log \csc (s-b) = 0.0977 \\
 & \quad \underline{2) 9.1757} \\
 \log \tan \frac{1}{2} A = 9.6419 & \log \tan \frac{1}{2} B = 9.5878 \\
 \frac{1}{2} A = 23^\circ 40.6' & \frac{1}{2} B = 21^\circ 9.7' \\
 \therefore A = 47^\circ 21.2' & \therefore B = 42^\circ 19.4'.
 \end{array}$$

$$\log \sin (s-a) = 9.8482$$

$$\log \sin (s-b) = 9.9023$$

$$\log \csc s = 0.0198$$

$$\log \csc (s-c) = 0.7900$$

$$\begin{array}{r} 2)0.5603 \\ \hline \end{array}$$

$$\log \tan \frac{1}{2}C = 0.2801$$

$$\frac{1}{2}C = 62^{\circ} 19'$$

$$\therefore C = 124^{\circ} 38'.$$

Art. 170.—Page 128.

2. Here $S = 109^{\circ}$, $S - A = 34^{\circ}$, $S - B = 27^{\circ}$, $S - C = 48^{\circ}$.

$$\log \cos S = 9.5126$$

$$\log \cos (S - A) = 9.9186$$

$$\log \sec (S - B) = 0.0501$$

$$\log \sec (S - C) = 0.1745$$

$$\begin{array}{r} 2)9.6558 \\ \hline \end{array}$$

$$\log \tan \frac{1}{2}a = 9.8279$$

$$\frac{1}{2}a = 33^{\circ} 55.9'$$

$$\therefore a = 67^{\circ} 51.8'.$$

$$\log \cos S = 9.5126$$

$$\log \cos (S - B) = 9.9499$$

$$\log \sec (S - C) = 0.1745$$

$$\log \sec (S - A) = 0.0814$$

$$\begin{array}{r} 2)9.7184 \\ \hline \end{array}$$

$$\log \tan \frac{1}{2}b = 9.8592$$

$$\frac{1}{2}b = 35^{\circ} 52.2'$$

$$\therefore b = 71^{\circ} 44.4'.$$

$$\log \cos S = 9.5126$$

$$\log \cos (S - C) = 9.8255$$

$$\log \sec (S - A) = 0.0814$$

$$\log \sec (S - B) = 0.0501$$

$$\begin{array}{r} 2)9.4696 \\ \hline \end{array}$$

$$\log \tan \frac{1}{2}c = 9.7348$$

$$\frac{1}{2}c = 28^{\circ} 30'$$

$$\therefore c = 57^{\circ}.$$

3. Here $S = 165^{\circ}$, $S - A = 45^{\circ}$, $S - B = 35^{\circ}$, $S - C = 85^{\circ}$.

$$\log \cos S = 9.9849$$

$$\log \cos (S - A) = 9.8495$$

$$\log \sec (S - B) = 0.0866$$

$$\log \sec (S - C) = 1.0597$$

$$\begin{array}{r} 2)0.9807 \\ \hline \end{array}$$

$$\log \tan \frac{1}{2}a = 0.4903$$

$$\frac{1}{2}a = 72^{\circ} 4.9'$$

$$\therefore a = 144^{\circ} 9.8'.$$

$$\log \cos S = 9.9849$$

$$\log \cos (S - B) = 9.9134$$

$$\log \sec (S - C) = 1.0597$$

$$\log \sec (S - A) = 0.1505$$

$$\begin{array}{r} 2)1.1085 \\ \hline \end{array}$$

$$\log \tan \frac{1}{2}b = 0.5542$$

$$\frac{1}{2}b = 74^{\circ} 24.3'$$

$$\therefore b = 148^{\circ} 48.6'.$$

$$\begin{aligned}
 \log \cos S &= 9.9849 \\
 \log \cos (S - C) &= 8.9403 \\
 \log \sec (S - A) &= 0.1505 \\
 \log \sec (S - B) &= 0.0866 \\
 &\quad \underline{2)9.1623} \\
 \log \tan \frac{1}{2} c &= 9.5811 \\
 \frac{1}{2} c &= 20^\circ 51.8' \\
 \therefore c &= 41^\circ 43.6'.
 \end{aligned}$$

4. Here $S = 124^\circ 40'$, $S - A = 33^\circ 30'$, $S - B = 39^\circ$, $S - C = 52^\circ 10'$.

$$\begin{array}{ll}
 \log \cos S &= 9.7550 & \log \cos S &= 9.7550 \\
 \log \cos (S - A) &= 9.9211 & \log \cos (S - B) &= 9.8905 \\
 \log \sec (S - B) &= 0.1095 & \log \sec (S - C) &= 0.2123 \\
 \log \sec (S - C) &= 0.2123 & \log \sec (S - A) &= 0.0789 \\
 &\quad \underline{2)9.9979} & &\quad \underline{2)9.9367} \\
 \log \tan \frac{1}{2} a &= 9.9989 & \log \tan \frac{1}{2} b &= 9.9683 \\
 \frac{1}{2} a &= 44^\circ 55.6' & \frac{1}{2} b &= 42^\circ 54.6' \\
 \therefore a &= 89^\circ 51.2' & \therefore b &= 85^\circ 49.2'.
 \end{array}$$

$$\begin{aligned}
 \log \cos S &= 9.7550 \\
 \log \cos (S - C) &= 9.7877 \\
 \log \sec (S - A) &= 0.0789 \\
 \log \sec (S - B) &= 0.1095 \\
 &\quad \underline{2)9.7311} \\
 \log \tan \frac{1}{2} c &= 9.8655 \\
 \frac{1}{2} c &= 36^\circ 15.9' \\
 \therefore c &= 72^\circ 31.8'.
 \end{aligned}$$

5. Here

$S = 102^\circ 40'$, $S - A = -35^\circ 36'$, $S - B = 71^\circ 29'$, $S - C = 66^\circ 47'$.

$$\begin{array}{ll}
 \log \cos S &= 9.3410 & \log \cos S &= 9.3410 \\
 \log \cos (S - A) &= 9.9102 & \log \cos (S - B) &= 9.5019 \\
 \log \sec (S - B) &= 0.4981 & \log \sec (S - C) &= 0.4043 \\
 \log \sec (S - C) &= 0.4043 & \log \sec (S - A) &= 0.0898 \\
 &\quad \underline{2)0.1536} & &\quad \underline{2)9.3370} \\
 \log \tan \frac{1}{2} a &= 0.0768 & \log \tan \frac{1}{2} b &= 9.6685 \\
 \frac{1}{2} a &= 50^\circ 2.3' & \frac{1}{2} b &= 24^\circ 59.4' \\
 \therefore a &= 100^\circ 4.6' & \therefore b &= 49^\circ 58.8'.
 \end{array}$$

$$\log \cos S = 9.3410$$

$$\log \cos (S - C) = 9.5957$$

$$\log \sec (S - A) = 0.0898$$

$$\log \sec (S - B) = 0.4981$$

$$\underline{2) 9.5246}$$

$$\log \tan \frac{1}{2} c = 9.7623$$

$$\frac{1}{2} c = 30^{\circ} 3'$$

$$\therefore c = 60^{\circ} 6'.$$

Art. 171. — Page 131.

$$4. \quad \sin C = \frac{\sin c \sin B}{\sin b}.$$

$$\log \sin c = 9.9549$$

$$\log \csc b = 0.0062$$

$$\log \sin B = 9.9979$$

$$\log \sin C = 9.9590$$

$$\therefore C = 65^{\circ} 30'.$$

$$\frac{1}{2} (b + c) = 82^{\circ},$$

$$\frac{1}{2} (b - c) = 17^{\circ} 40'.$$

$$\frac{1}{2} (B + C) = 80^{\circ} 35',$$

$$\frac{1}{2} (B - C) = 15^{\circ} 5'.$$

$$\cot \frac{1}{2} A = \frac{\sin \frac{1}{2} (b + c)}{\sin \frac{1}{2} (b - c)} \tan \frac{1}{2} (B - C).$$

$$\log \sin \frac{1}{2} (b + c) = 9.9958$$

$$\log \csc \frac{1}{2} (b - c) = 0.5179$$

$$\log \tan \frac{1}{2} (B - C) = \underline{9.4306}$$

$$\log \cot \frac{1}{2} A = \underline{9.9443}$$

$$\frac{1}{2} A = 48^{\circ} 40'$$

$$\therefore A = 97^{\circ} 20'.$$

$$\tan \frac{1}{2} a = \frac{\sin \frac{1}{2} (B + C)}{\sin \frac{1}{2} (B - C)} \tan \frac{1}{2} (b - c).$$

$$\log \sin \frac{1}{2} (B + C) = 9.9941$$

$$\log \csc \frac{1}{2} (B - C) = 0.5847$$

$$\log \tan \frac{1}{2} (b - c) = \underline{9.5031}$$

$$\log \tan \frac{1}{2} a = \underline{0.0819}$$

$$\frac{1}{2} a = 50^{\circ} 22.3'$$

$$\therefore a = 100^{\circ} 44.6'.$$

$$5. \quad \sin B = \frac{\sin b \sin A}{\sin a}.$$

$$\log \sin b = 9.9446$$

$$\log \csc a = 0.1919$$

$$\log \sin A = 9.6946$$

$$\log \sin B = 9.8311$$

$$\therefore B = 42^{\circ} 40', \text{ or } 137^{\circ} 20'.$$

$$\frac{1}{2} (b + a) = 79^{\circ} 10',$$

$$\frac{1}{2} (b - a) = 39^{\circ} 10'.$$

$$\frac{1}{2} (B + A) = 36^{\circ} 10', \text{ or } 83^{\circ} 30'.$$

$$\frac{1}{2} (B - A) = 6^{\circ} 30', \text{ or } 53^{\circ} 50'.$$

$$\cot \frac{1}{2} C = \frac{\sin \frac{1}{2} (b + a)}{\sin \frac{1}{2} (b - a)} \tan \frac{1}{2} (B - A).$$

$$\tan \frac{1}{2} c = \frac{\sin \frac{1}{2} (B + A)}{\sin \frac{1}{2} (B - A)} \tan \frac{1}{2} (b - a).$$

Using the first value of B , we have :

$$\log \sin \frac{1}{2} (b + a) = 9.9922$$

$$\log \csc \frac{1}{2} (b - a) = 0.1996$$

$$\log \tan \frac{1}{2} (B - A) = \underline{9.0567}$$

$$\log \cot \frac{1}{2} C = \underline{9.2485}$$

$$\frac{1}{2} C = 79^{\circ} 57'$$

$$\therefore C = 159^{\circ} 54'.$$

$$\log \sin \frac{1}{2} (B + A) = 9.7710$$

$$\log \csc \frac{1}{2} (B - A) = 0.9461$$

$$\log \tan \frac{1}{2} (b - a) = \underline{9.9110}$$

$$\log \tan \frac{1}{2} c = \underline{0.6281}$$

$$\frac{1}{2} c = 76^{\circ} 45.1'$$

$$\therefore c = 153^{\circ} 30.2'.$$

Using the second value of B , we have:

$$\begin{array}{ll}
 \log \sin \frac{1}{2}(b+a) = 9.9922 & \log \sin \frac{1}{2}(B+A) = 9.9972 \\
 \log \csc \frac{1}{2}(b-a) = 0.1996 & \log \csc \frac{1}{2}(B-A) = 0.0930 \\
 \log \tan \frac{1}{2}(B-A) = 0.1361 & \log \tan \frac{1}{2}(b-a) = 9.9110 \\
 \log \cot \frac{1}{2}C = 0.3279 & \log \tan \frac{1}{2}c = 0.0012 \\
 \frac{1}{2}C = 25^{\circ} 10.3' & \frac{1}{2}c = 45^{\circ} 4.8' \\
 \therefore C = 50^{\circ} 20.6' & \therefore c = 90^{\circ} 9.6'.
 \end{array}$$

$$6. \quad \sin A = \frac{\sin a \sin C}{\sin c}.$$

$$\begin{array}{l}
 \log \sin a = 9.9561 \\
 \log \csc c = 0.2562 \\
 \log \sin C = 9.7973 \\
 \log \sin A = 0.0096
 \end{array}$$

Since $\log \sin A$ is *positive*, the triangle is impossible.

$$7. \quad \sin C = \frac{\sin c \sin A}{\sin a}.$$

$$\begin{array}{l}
 \log \sin c = 9.9958 \\
 \log \csc a = 0.0252 \\
 \log \sin A = 9.9790 \\
 \log \sin C = 0.0000 \\
 \therefore C = 90^{\circ}.
 \end{array}$$

$$\cos B = \frac{\tan a}{\tan c}.$$

$$\begin{array}{l}
 + \\
 \log \tan a = 0.4549 \\
 \log \tan c = 0.8522 \\
 \log \cos B = 9.6027 \\
 180^{\circ} - B = 66^{\circ} 23.1' \\
 \therefore B = 113^{\circ} 36.9'.
 \end{array}$$

$$\cos b = \frac{\cos c}{\cos a}.$$

$$\begin{array}{l}
 + \\
 \log \cos c = 9.1436 \\
 \log \cos a = 9.5199 \\
 \log \cos b = 9.6237 \\
 180^{\circ} - b = 65^{\circ} 8.1' \\
 \therefore b = 114^{\circ} 51.9'.
 \end{array}$$

$$8. \quad \sin B = \frac{\sin b \sin C}{\sin c}.$$

$$\begin{array}{l}
 \log \sin b = 9.9770 \\
 \log \csc c = 0.1845 \\
 \log \sin C = 9.8066 \\
 \log \sin B = 9.9681
 \end{array}$$

$$\begin{array}{l}
 \frac{1}{2}(b+c) = 74^{\circ} 40', \\
 \frac{1}{2}(b-c) = 33^{\circ} 50'.
 \end{array}$$

$$\begin{array}{l}
 \frac{1}{2}(B+C) = 54^{\circ} 4', \text{ or } 75^{\circ} 46'. \\
 \frac{1}{2}(B-C) = 14^{\circ} 14', \text{ or } 35^{\circ} 56'.
 \end{array}$$

$$\therefore B = 68^{\circ} 18', \text{ or } 111^{\circ} 42'.$$

$$\cot \frac{1}{2}A = \frac{\sin \frac{1}{2}(b+c)}{\sin \frac{1}{2}(b-c)} \tan \frac{1}{2}(B-C). \quad \tan \frac{1}{2}a = \frac{\sin \frac{1}{2}(B+C)}{\sin \frac{1}{2}(B-C)} \tan \frac{1}{2}(b-c).$$

Using the first value of B , we have :

$$\begin{array}{ll}
 \log \sin \frac{1}{2}(b+c) = 9.9843 & \log \sin \frac{1}{2}(B+C) = 9.9084 \\
 \log \csc \frac{1}{2}(b-c) = 0.2543 & \log \csc \frac{1}{2}(B-C) = 0.6093 \\
 \log \tan \frac{1}{2}(B-C) = 9.4042 & \log \tan \frac{1}{2}(b-c) = 9.8263 \\
 \log \cot \frac{1}{2}A = 9.6428 & \log \tan \frac{1}{2}a = 0.3440 \\
 \frac{1}{2}A = 66^{\circ} 16.9' & \frac{1}{2}a = 65^{\circ} 37.9' \\
 \therefore A = 132^{\circ} 33.8'. & \therefore a = 131^{\circ} 15.8'.
 \end{array}$$

Using the second value of B , we have :

$$\begin{array}{ll}
 \log \sin \frac{1}{2}(b+c) = 9.9843 & \log \sin \frac{1}{2}(B+C) = 9.9865 \\
 \log \csc \frac{1}{2}(b-c) = 0.2543 & \log \csc \frac{1}{2}(B-C) = 0.2315 \\
 \log \tan \frac{1}{2}(B-C) = 9.8602 & \log \tan \frac{1}{2}(b-c) = 9.8263 \\
 \log \cot \frac{1}{2}A = 0.0988 & \log \tan \frac{1}{2}a = 0.0443 \\
 \frac{1}{2}A = 38^{\circ} 32.3' & \frac{1}{2}a = 47^{\circ} 55' \\
 \therefore A = 77^{\circ} 4.6'. & \therefore a = 95^{\circ} 50'.
 \end{array}$$

$$9. \quad \sin A = \frac{\sin a \sin B}{\sin b}$$

$$\begin{array}{l}
 \log \sin a = 9.4821 \\
 \log \csc b = 0.5686 \\
 \log \sin B = 9.9134 \\
 \log \sin A = 9.9641 \\
 \therefore A = 67^{\circ} 1.7', \text{ or } 112^{\circ} 58.3'.
 \end{array}$$

Since both values of A are $< B$, while a is given $> b$, the triangle is impossible.

$$\begin{array}{ll}
 10. \quad \sin C = \frac{\sin c \sin A}{\sin a} & \frac{1}{2}(c+a) = 96^{\circ} 35', \\
 \log \sin c = 9.8241 & \frac{1}{2}(c-a) = 41^{\circ} 35'. \\
 \log \csc a = 0.0866 & \frac{1}{2}(C+A) = 94^{\circ} 33.95', \\
 \log \sin A = 9.8297 & \frac{1}{2}(C-A) = 52^{\circ} 3.95'. \\
 \log \sin C = 9.7404 & \\
 \therefore C = 146^{\circ} 37.9'. &
 \end{array}$$

$$\begin{array}{ll}
 \cot \frac{1}{2}B = \frac{\sin \frac{1}{2}(c+a)}{\sin \frac{1}{2}(c-a)} \tan \frac{1}{2}(C-A). & \tan \frac{1}{2}b = \frac{\sin \frac{1}{2}(C+A)}{\sin \frac{1}{2}(C-A)} \tan \frac{1}{2}(c-a). \\
 \log \sin \frac{1}{2}(c+a) = 9.9972 & \log \sin \frac{1}{2}(C+A) = 9.9987 \\
 \log \csc \frac{1}{2}(c-a) = 0.1780 & \log \csc \frac{1}{2}(C-A) = 0.1031 \\
 \log \tan \frac{1}{2}(C-A) = 0.1082 & \log \tan \frac{1}{2}(c-a) = 9.9481 \\
 \log \cot \frac{1}{2}B = 0.2834 & \log \tan \frac{1}{2}b = 0.0499 \\
 \frac{1}{2}B = 27^{\circ} 30.3' & \frac{1}{2}b = 48^{\circ} 17.2' \\
 \therefore B = 55^{\circ} 0.6'. & \therefore b = 96^{\circ} 34.4'.
 \end{array}$$

Art. 172. — Page 133.

$$\begin{array}{ll}
2. & \sin b = \frac{\sin B \sin c}{\sin C}. & \frac{1}{2}(B + C) = 98^\circ, \\
& \log \sin B = 9.9537 & \frac{1}{2}(B - C) = 18^\circ. \\
& \log \csc C = 0.0066 & \frac{1}{2}(b + c) = 99^\circ 25', \\
& \log \sin c = 9.9976 & \frac{1}{2}(b - c) = 15^\circ 25'. \\
& \log \sin b = 9.9579 & \\
& \therefore b = 114^\circ 50'. & \\
\cot \frac{1}{2} A = \frac{\sin \frac{1}{2}(b + c)}{\sin \frac{1}{2}(b - c)} \tan \frac{1}{2}(B - C). & \tan \frac{1}{2} a = \frac{\sin \frac{1}{2}(B + C)}{\sin \frac{1}{2}(B - C)} \tan \frac{1}{2}(b - c). \\
\log \sin \frac{1}{2}(b + c) = 9.9941 & \log \sin \frac{1}{2}(B + C) = 9.9958 \\
\log \csc \frac{1}{2}(b - c) = 0.5754 & \log \csc \frac{1}{2}(B - C) = 0.5100 \\
\log \tan \frac{1}{2}(B - C) = 9.5118 & \log \tan \frac{1}{2}(b - c) = 9.4405 \\
\log \cot \frac{1}{2} A = 0.0813 & \log \tan \frac{1}{2} a = 9.9463 \\
\frac{1}{2} A = 39^\circ 40' & \frac{1}{2} a = 41^\circ 28' \\
\therefore A = 79^\circ 20'. & \therefore a = 82^\circ 56'.
\end{array}$$

$$\begin{array}{ll}
3. & \sin a = \frac{\sin A \sin b}{\sin B}. & \frac{1}{2}(B + A) = 136^\circ, \\
& \log \sin A = 9.8711 & \frac{1}{2}(B - A) = 4^\circ. \\
& \log \csc B = 0.1919 & \frac{1}{2}(b + a) = 97^\circ 12', \text{ or } 119^\circ 48'. \\
& \log \sin b = 9.9023 & \frac{1}{2}(b - a) = 29^\circ 48', \text{ or } 7^\circ 12'. \\
& \log \sin a = 9.9653 & \\
& \therefore a = 67^\circ 24', \text{ or } 112^\circ 36'. & \\
\cot \frac{1}{2} C = \frac{\sin \frac{1}{2}(b + a)}{\sin \frac{1}{2}(b - a)} \tan \frac{1}{2}(B - A). & \tan \frac{1}{2} c = \frac{\sin \frac{1}{2}(B + A)}{\sin \frac{1}{2}(B - A)} \tan \frac{1}{2}(b - a).
\end{array}$$

Using the first value of a , we have :

$$\begin{array}{ll}
\log \sin \frac{1}{2}(b + a) = 9.9966 & \log \sin \frac{1}{2}(B + A) = 9.8418 \\
\log \csc \frac{1}{2}(b - a) = 0.3036 & \log \csc \frac{1}{2}(B - A) = 1.1564 \\
\log \tan \frac{1}{2}(B - A) = 8.8446 & \log \tan \frac{1}{2}(b - a) = 9.7579 \\
\log \cot \frac{1}{2} C = 9.1448 & \log \tan \frac{1}{2} c = 0.7561 \\
\frac{1}{2} C = 82^\circ 3.2' & \frac{1}{2} c = 80^\circ 3.2' \\
\therefore C = 164^\circ 6.4'. & \therefore c = 160^\circ 6.4'.
\end{array}$$

Using the second value of a , we have :

$$\begin{array}{ll}
\log \sin \frac{1}{2}(b + a) = 9.9384 & \log \sin \frac{1}{2}(B + A) = 9.8418 \\
\log \csc \frac{1}{2}(b - a) = 0.9019 & \log \csc \frac{1}{2}(B - A) = 1.1564 \\
\log \tan \frac{1}{2}(B - A) = 8.8446 & \log \tan \frac{1}{2}(b - a) = 9.1015 \\
\log \cot \frac{1}{2} C = 9.6849 & \log \tan \frac{1}{2} c = 0.0997 \\
\frac{1}{2} C = 64^\circ 10.3' & \frac{1}{2} c = 51^\circ 31.2' \\
\therefore C = 128^\circ 20.6'. & \therefore c = 103^\circ 2.4'.
\end{array}$$

$$4. \quad \sin c = \frac{\sin C \sin a}{\sin A}.$$

$$\log \sin C = 9.9904$$

$$\log \csc A = 0.0541$$

$$\log \sin a = \underline{9.9555}$$

$$\log \sin c = \underline{0.0000}$$

$$\therefore c = 90^\circ.$$

The triangle is a quadrantal triangle, and the sides a' and c' of its polar triangle are 118° and 78° .

$$\overset{-}{\cos b'} = \frac{\overset{+}{\cos c'}}{\overset{-}{\cos a'}}$$

$$\overset{-}{\cos B'} = \frac{\overset{-}{\tan a'}}{\overset{+}{\tan c'}}$$

$$\log \cos c' = 9.3179$$

$$\log \tan a' = 0.2743$$

$$\log \cos a' = \underline{9.6716}$$

$$\log \tan c' = \underline{0.6725}$$

$$\log \cos b' = \underline{9.6463}$$

$$\log \cos B' = \underline{9.6018}$$

$$\therefore 180^\circ - b' = 63^\circ 42.7'.$$

$$\therefore 180^\circ - B' = 66^\circ 26.2'.$$

Therefore in the given triangle, $B = 63^\circ 42.7'$, and $b = 66^\circ 26.2'$.

$$5. \quad \sin b = \frac{\sin B \sin a}{\sin A}.$$

$$\log \sin B = 9.9624$$

$$\log \csc A = 0.1418$$

$$\log \sin a = \underline{9.9948}$$

$$\log \sin b = \underline{0.0990}$$

Since $\log \sin b$ is positive, the triangle is impossible.

$$6. \quad \sin b = \frac{\sin B \sin c}{\sin C}.$$

$$\frac{1}{2}(C + B) = 84^\circ 30',$$

$$\frac{1}{2}(C - B) = 62^\circ 10'.$$

$$\log \sin B = 9.5798$$

$$\frac{1}{2}(c + b) = 82^\circ 51.05',$$

$$\log \csc C = 0.2600$$

$$\frac{1}{2}(c - b) = 55^\circ 28.95'.$$

$$\log \sin c = \underline{9.8227}$$

$$\log \sin b = \underline{9.6625}$$

$$\therefore b = 27^\circ 22.1'.$$

$$\cot \frac{1}{2} A = \frac{\sin \frac{1}{2}(c + b)}{\sin \frac{1}{2}(c - b)} \tan \frac{1}{2}(C - B). \quad \tan \frac{1}{2} a = \frac{\sin \frac{1}{2}(C + B)}{\sin \frac{1}{2}(C - B)} \tan \frac{1}{2}(c - b).$$

$$\log \sin \frac{1}{2}(c + b) = 9.9966$$

$$\log \sin \frac{1}{2}(C + B) = 9.9980$$

$$\log \csc \frac{1}{2}(c - b) = 0.0841$$

$$\log \csc \frac{1}{2}(C - B) = 0.0534$$

$$\log \tan \frac{1}{2}(C - B) = \underline{0.2774}$$

$$\log \tan \frac{1}{2}(c - b) = \underline{0.1626}$$

$$\log \cot \frac{1}{2} A = \underline{0.3581}$$

$$\log \tan \frac{1}{2} a = \underline{0.2140}$$

$$\frac{1}{2} A = 23^\circ 40.6'$$

$$\frac{1}{2} a = 58^\circ 34.6'$$

$$\therefore A = 47^\circ 21.2'.$$

$$\therefore a = 117^\circ 9.2'.$$

$$7. \quad \sin a = \frac{\sin A \sin c}{\sin C}.$$

$$\log \sin A = 9.9446$$

$$\log \csc C = 0.1950$$

$$\log \sin c = \underline{9.6946}$$

$$\log \sin a = \underline{9.8342}$$

$$\therefore a = 43^\circ 3.1', \text{ or } 136^\circ 56.9'.$$

$$\frac{1}{2}(C + A) = 101^\circ,$$

$$\frac{1}{2}(C - A) = 39^\circ 20'.$$

$$\frac{1}{2}(c + a) = 96^\circ 41.55',$$

$$\text{or } 143^\circ 38.45'.$$

$$\frac{1}{2}(c - a) = 53^\circ 38.45',$$

$$\text{or } 6^\circ 41.55'.$$

$$\cot \frac{1}{2}B = \frac{\sin \frac{1}{2}(c + a)}{\sin \frac{1}{2}(c - a)} \tan \frac{1}{2}(C - A). \quad \tan \frac{1}{2}b = \frac{\sin \frac{1}{2}(C + A)}{\sin \frac{1}{2}(C - A)} \tan \frac{1}{2}(c - a).$$

Using the first value of a , we have :

$$\log \sin \frac{1}{2}(c + a) = 9.9971$$

$$\log \csc \frac{1}{2}(c - a) = 0.0940$$

$$\log \tan \frac{1}{2}(C - A) = \underline{9.9135}$$

$$\log \cot \frac{1}{2}B = 0.0046$$

$$\frac{1}{2}B = 44^\circ 41.9'$$

$$\therefore B = 89^\circ 23.8'.$$

$$\log \sin \frac{1}{2}(C + A) = 9.9919$$

$$\log \csc \frac{1}{2}(C - A) = 0.1980$$

$$\log \tan \frac{1}{2}(c - a) = \underline{0.1330}$$

$$\log \tan \frac{1}{2}b = 0.3229$$

$$\frac{1}{2}b = 64^\circ 34.2'$$

$$\therefore b = 129^\circ 8.4'.$$

Using the second value of a , we have :

$$\log \sin \frac{1}{2}(c + a) = 9.7730$$

$$\log \csc \frac{1}{2}(c - a) = 0.9335$$

$$\log \tan \frac{1}{2}(C - A) = \underline{9.9135}$$

$$\log \cot \frac{1}{2}B = 0.6200$$

$$\frac{1}{2}B = 13^\circ 29.3'$$

$$\therefore B = 26^\circ 58.6'.$$

$$\log \sin \frac{1}{2}(C + A) = 9.9919$$

$$\log \csc \frac{1}{2}(C - A) = 0.1980$$

$$\log \tan \frac{1}{2}(c - a) = \underline{9.0695}$$

$$\log \tan \frac{1}{2}b = 9.2594$$

$$\frac{1}{2}b = 10^\circ 17.9'$$

$$\therefore b = 20^\circ 35.8'.$$

$$8. \quad \sin c = \frac{\sin C \sin b}{\sin B}.$$

$$\log \sin C = 9.9950$$

$$\log \csc B = 0.0194$$

$$\log \sin b = \underline{9.9252}$$

$$\log \sin c = \underline{9.9396}$$

$$\therefore c = 60^\circ 28.6', \text{ or } 119^\circ 31.4'.$$

Since both values of c are $< b$, while C is given $> B$, the triangle is impossible.

Art. 173.—Pages 134 and 136.

1. Let B and G denote the positions of Boston and Greenwich, respectively, and P the north-pole.

Denote the arcs PG , PB , and BG by b , g , and p .

Then in the triangle PBG ,

$$\angle P = 71^\circ 4', b = 90^\circ - 51^\circ 29' = 38^\circ 31', g = 90^\circ - 42^\circ 21' = 47^\circ 39'.$$

By Art. 162, we have :

$$\tan \frac{1}{2}(G - B) = \frac{\sin \frac{1}{2}(g - b)}{\sin \frac{1}{2}(g + b)} \cot \frac{1}{2}P,$$

$$\tan \frac{1}{2}(G + B) = \frac{\cos \frac{1}{2}(g - b)}{\cos \frac{1}{2}(g + b)} \cot \frac{1}{2}P.$$

From the data,

$$\frac{1}{2}(g - b) = 4^\circ 34', \frac{1}{2}(g + b) = 43^\circ 5', \frac{1}{2}P = 35^\circ 32'.$$

$$\log \sin \frac{1}{2}(g - b) = 8.9009$$

$$\log \cos \frac{1}{2}(g - b) = 9.9987$$

$$\log \csc \frac{1}{2}(g + b) = 0.1656$$

$$\log \sec \frac{1}{2}(g + b) = 0.1365$$

$$\log \cot \frac{1}{2}P = 0.1462$$

$$\log \cot \frac{1}{2}P = 0.1462$$

$$\log \tan \frac{1}{2}(G - B) = 9.2127$$

$$\log \tan \frac{1}{2}(G + B) = 0.2814$$

$$\therefore \frac{1}{2}(G - B) = 9^\circ 16.1'.$$

$$\therefore \frac{1}{2}(G + B) = 62^\circ 23.2'.$$

$$\therefore B = \frac{1}{2}(G + B) - \frac{1}{2}(G - B) = 53^\circ 7.1',$$

and

$$G = \frac{1}{2}(G + B) + \frac{1}{2}(G - B) = 71^\circ 39.3'.$$

$$\tan \frac{1}{2}p = \frac{\sin \frac{1}{2}(G + B)}{\sin \frac{1}{2}(G - B)} \tan \frac{1}{2}(g - b).$$

$$\log \sin \frac{1}{2}(G + B) = 9.9475$$

$$\log \csc \frac{1}{2}(G - B) = 0.7930$$

$$\log \tan \frac{1}{2}(g - b) = 8.9023$$

$$\log \tan \frac{1}{2}p = 9.6428$$

$$\frac{1}{2}p = 23^\circ 43.1'$$

$$\therefore p = 47^\circ 26.2'.$$

$$47^\circ 26.2' = 2846.2'.$$

$$360^\circ = 21600'.$$

$$\text{Circumference of earth} = 7912 \times 3.1416.$$

$$\therefore p \text{ (in miles)} = \frac{2846.2}{21600} \times 7912 \times 3.1416.$$

$$\log 2846.2 = 3.4542$$

$$\text{colog } 21600 = 5.6655$$

$$\log 7912 = 3.8983$$

$$\log 3.1416 = 0.4971$$

$$\log p = 3.5151$$

$$\therefore p = 3274.3.$$

Therefore the shortest distance between the places is 3274.3 miles; the bearing of Greenwich from Boston is N. $53^\circ 7.1'$ E., and of Boston from Greenwich, N. $71^\circ 39.3'$ W.

2. Let C and V denote the positions of Calcutta and Valparaiso, respectively, and P the north-pole.

Denote the arcs PV , PC , and CV by c , v , and p .

Then in the triangle PCV ,

$$\angle P = 88^\circ 19' + 71^\circ 42' = 160^\circ 1', \quad c = 90^\circ + 33^\circ 2' = 123^\circ 2',$$

$$v = 90^\circ - 22^\circ 33' = 67^\circ 27'.$$

By Art. 162, we have :

$$\tan \frac{1}{2}(C - V) = \frac{\sin \frac{1}{2}(c - v)}{\sin \frac{1}{2}(c + v)} \cot \frac{1}{2}P,$$

$$\tan \frac{1}{2}(C + V) = \frac{\cos \frac{1}{2}(c - v)}{\cos \frac{1}{2}(c + v)} \cot \frac{1}{2}P.$$

From the data, $\frac{1}{2}(c - v) = 27^\circ 47.5'$, $\frac{1}{2}(c + v) = 95^\circ 14.5'$, $\frac{1}{2}P = 80^\circ 0.5'$.

$$\log \sin \frac{1}{2}(c - v) = 9.6686$$

$$\log \cos \frac{1}{2}(c - v) = 9.9468$$

$$\log \csc \frac{1}{2}(c + v) = 0.0018$$

$$\log \sec \frac{1}{2}(c + v) = 1.0393$$

$$\log \cot \frac{1}{2}P = 9.2459$$

$$\log \cot \frac{1}{2}P = 9.2459$$

$$\log \tan \frac{1}{2}(C - V) = 8.9163$$

$$\log \tan \frac{1}{2}(C + V) = 0.2320$$

$$\therefore \frac{1}{2}(C - V) = 4^\circ 42.9'.$$

$$180^\circ - \frac{1}{2}(C + V) = 59^\circ 37.5'$$

$$\therefore \frac{1}{2}(C + V) = 120^\circ 22.5'.$$

$$\therefore C = \frac{1}{2}(C + V) + \frac{1}{2}(C - V) = 125^\circ 5.4',$$

and

$$V = \frac{1}{2}(C + V) - \frac{1}{2}(C - V) = 115^\circ 39.6'.$$

$$\tan \frac{1}{2}p = \frac{\sin \frac{1}{2}(C + V)}{\sin \frac{1}{2}(C - V)} \tan \frac{1}{2}(c - v).$$

$$\log \sin \frac{1}{2}(C + V) = 9.9359$$

$$\log \csc \frac{1}{2}(C - V) = 1.0852$$

$$\log \tan \frac{1}{2}(c - v) = 9.7218$$

$$\log \tan \frac{1}{2}p = 0.7429$$

$$\frac{1}{2}p = 79^\circ 45.2'$$

$$\therefore p = 159^\circ 30.4'.$$

$$159^\circ 30.4' = 9570.4'.$$

$$360^\circ = 21600'.$$

Circumference of earth = 7912×3.1416 .

$$\therefore p \text{ (in miles)} = \frac{9570.4}{21600} \times 7912 \times 3.1416.$$

$$\log 9570.4 = 3.9809$$

$$\text{colog } 21600 = 5.6655$$

$$\log 7912 = 3.8983$$

$$\log 3.1416 = 0.4971$$

$$\log p = 4.0418$$

$$\therefore p = 11010.$$

Therefore the shortest distance between the places is 11010 miles; the bearing of Calcutta from Valparaiso is N. $115^{\circ} 39.6'$ E., or S. $64^{\circ} 20.4'$ E., and of Valparaiso from Calcutta is N. $125^{\circ} 54.4'$ W., or S. $54^{\circ} 54.6'$ W.

3. Let S and Q denote the positions of Sandy Hook and Queens-town, respectively, P the north pole, and X the intersection of the arc QS with the meridian of 50° W.

Denote the arcs PQ and PS by s and q .

Then in the triangle PQS ,

$$\begin{aligned}\angle P &= 74^{\circ} 1' - 8^{\circ} 19' = 65^{\circ} 42', & q &= 90^{\circ} - 40^{\circ} 28' = 49^{\circ} 32', \\ s &= 90^{\circ} - 51^{\circ} 50' = 38^{\circ} 10' .\end{aligned}$$

By Art. 162, we have :

$$\tan \frac{1}{2} (Q - S) = \frac{\sin \frac{1}{2} (q - s)}{\sin \frac{1}{2} (q + s)} \cot \frac{1}{2} P,$$

$$\tan \frac{1}{2} (Q + S) = \frac{\cos \frac{1}{2} (q - s)}{\cos \frac{1}{2} (q + s)} \cot \frac{1}{2} P.$$

From the data,

$$\frac{1}{2} (q - s) = 5^{\circ} 41', \quad \frac{1}{2} (q + s) = 43^{\circ} 51', \quad \frac{1}{2} P = 32^{\circ} 51'.$$

$\log \sin \frac{1}{2} (q - s) = 8.9957$	$\log \cos \frac{1}{2} (q - s) = 9.9979$
$\log \csc \frac{1}{2} (q + s) = 0.1594$	$\log \sec \frac{1}{2} (q + s) = 0.1419$
$\log \cot \frac{1}{2} P = 0.1900$	$\log \cot \frac{1}{2} P = 0.1900$

$\log \tan \frac{1}{2} (Q - S) = 9.3451$	$\log \tan \frac{1}{2} (Q + S) = 0.3298$
--	--

$$\therefore \frac{1}{2} (Q - S) = 12^{\circ} 28.9'.$$

$$\therefore \frac{1}{2} (Q + S) = 64^{\circ} 55.5'.$$

Then in the triangle PQX ,

$$\angle P = 50^{\circ} - 8^{\circ} 19' = 41^{\circ} 41',$$

$$\angle Q = \frac{1}{2} (Q + S) + \frac{1}{2} (Q - S) = 77^{\circ} 24.4', \quad PQ = 38^{\circ} 10'.$$

By Arts. 160 and 161, we have :

$$\tan \frac{1}{2} (PX - QX) = \frac{\sin \frac{1}{2} (Q - QPX)}{\sin \frac{1}{2} (Q + QPX)} \tan \frac{1}{2} PQ.$$

$$\tan \frac{1}{2} (PX + QX) = \frac{\cos \frac{1}{2} (Q - QPX)}{\cos \frac{1}{2} (Q + QPX)} \tan \frac{1}{2} PQ.$$

From the data,

$$\frac{1}{2} (Q - QPX) = 17^{\circ} 51.7', \quad \frac{1}{2} (Q + QPX) = 59^{\circ} 32.7', \quad \frac{1}{2} PQ = 19^{\circ} 5'.$$

$\log \sin \frac{1}{2} (Q - QPX) = 9.4868$	$\log \cos \frac{1}{2} (Q - QPX) = 9.9785$
$\log \csc \frac{1}{2} (Q + QPX) = 0.0645$	$\log \sec \frac{1}{2} (Q + QPX) = 0.2951$
$\log \tan \frac{1}{2} PQ = 9.5390$	$\log \tan \frac{1}{2} PQ = 9.5390$

$\log \tan \frac{1}{2} (PX - QX) = 9.0903$	$\log \tan \frac{1}{2} (PX + QX) = 9.8126$
--	--

$$\therefore \frac{1}{2} (PX - QX) = 7^{\circ} 1.2'.$$

$$\therefore \frac{1}{2} (PX + QX) = 33^{\circ} 0.4'.$$

$$\therefore PX = \frac{1}{2} (PX + QX) + \frac{1}{2} (PX - QX) = 40^{\circ} 1.6'.$$

Therefore the latitude of $X = 90^{\circ} - 40^{\circ} 1.6' = 49^{\circ} 58.4'$ N.

4. Denote the arcs SP , SZ , and PZ (see Fig., p. 135) by z , p , and s , and their sum by $2s'$.

Then in the triangle SPZ ,

$$z = 90^\circ - 18^\circ 36' = 71^\circ 24', \quad p = 90^\circ - 14^\circ 18' = 75^\circ 42',$$

$$s = 90^\circ - 50^\circ 13' = 39^\circ 47'.$$

By Art. 158,

$$\cos \frac{1}{2} P = \sqrt{\frac{\sin s' \sin (s' - p)}{\sin s \sin z}}.$$

From the data,

$$s' = 93^\circ 26.5', \quad s' - p = 17^\circ 44.5'.$$

$$\log \sin s' = 9.9992$$

$$\log \sin (s' - p) = 9.4839$$

$$\log \csc s = 0.1939$$

$$\log \csc z = 0.0233$$

$$\hline 2) 9.7003$$

$$\log \cos \frac{1}{2} P = 9.8501$$

$$\therefore \frac{1}{2} P = 44^\circ 55', \text{ and } P = 89^\circ 50'.$$

To $89^\circ 50'$ corresponds 5 h. 59 m. 20 s. of time.

Therefore the hour of the day is 6 h. 0 m. 40 s. A.M.

The difference between the local and Greenwich time is 2 h. 59 m. 20 s., which corresponds to $44^\circ 50'$ of longitude.

Hence the longitude of the vessel is $44^\circ 50' W$.

5. Denote the arcs SP , SZ , and PZ (see Fig., p. 135) by z , p , and s . Then in the triangle SPZ ,

$$z = 90^\circ + 12^\circ = 102^\circ, \quad s = 90^\circ - 37^\circ 48' = 52^\circ 12', \quad \angle P = 60^\circ.$$

By Art 162, we have :

$$\tan \frac{1}{2} (Z - S) = \frac{\sin \frac{1}{2} (z - s)}{\sin \frac{1}{2} (z + s)} \cot \frac{1}{2} P,$$

$$\tan \frac{1}{2} (Z + S) = \frac{\cos \frac{1}{2} (z - s)}{\cos \frac{1}{2} (z + s)} \cot \frac{1}{2} P.$$

From the data, $\frac{1}{2} (z - s) = 24^\circ 54'$, $\frac{1}{2} (z + s) = 77^\circ 6'$, $\frac{1}{2} P = 30^\circ$.

$$\log \sin \frac{1}{2} (z - s) = 9.6243$$

$$\log \cos \frac{1}{2} (z - s) = 9.9577$$

$$\log \csc \frac{1}{2} (z + s) = 0.0111$$

$$\log \sec \frac{1}{2} (z + s) = 0.6512$$

$$\log \cot \frac{1}{2} P = 0.2386$$

$$\log \cot \frac{1}{2} P = 0.2386$$

$$\log \tan \frac{1}{2} (Z - S) = 9.8740$$

$$\log \tan \frac{1}{2} (Z + S) = 0.8475$$

$$\therefore \frac{1}{2} (Z - S) = 36^\circ 48.1'.$$

$$\therefore \frac{1}{2} (Z + S) = 81^\circ 54.8'.$$

$$\begin{aligned}\tan \frac{1}{2} p &= \frac{\sin \frac{1}{2} (Z + S)}{\sin \frac{1}{2} (Z - S)} \tan \frac{1}{2} (z - s). \\ \log \sin \frac{1}{2} (Z + S) &= 9.9957 \\ \log \csc \frac{1}{2} (Z - S) &= 0.2225 \\ \log \tan \frac{1}{2} (z - s) &= \underline{9.6667} \\ \log \tan \frac{1}{2} p &= \underline{9.8849}\end{aligned}$$

$$\therefore \frac{1}{2} p = 37^{\circ} 29.6', \text{ and } p = 74^{\circ} 59.2'.$$

Therefore the altitude of the sun is $90^{\circ} - 74^{\circ} 59.2'$, or $15^{\circ} 0.8'$.

6. Denote the arcs SP , SZ , and PZ (see Fig., p. 135) by z , p , and s , and their sum by $2s'$.

Then in the triangle SPZ ,

$$z = 90^{\circ} + 3^{\circ} = 93^{\circ}, p = 90^{\circ} - 25^{\circ} 46' = 64^{\circ} 14', s = 90^{\circ} + 37^{\circ} 49' = 127^{\circ} 49'.$$

$$\text{By Art. 158,} \quad \cos \frac{1}{2} P = \sqrt{\frac{\sin s' \sin (s' - p)}{\sin s \sin z}}.$$

From the data, $s' = 142^{\circ} 31.5'$, $s' - p = 78^{\circ} 17.5'$.

$$\begin{aligned}\log \sin s' &= 9.7842 \\ \log \sin (s' - p) &= 9.9908 \\ \log \csc s &= 0.1024 \\ \log \csc z &= \underline{0.0006} \\ &2) 9.8780 \\ \log \cos \frac{1}{2} P &= \underline{9.9390}\end{aligned}$$

$$\therefore \frac{1}{2} P = 29^{\circ} 40', \text{ and } P = 59^{\circ} 20'.$$

To $59^{\circ} 20'$ corresponds 3 h. 57 m. 20 s. of time.

Therefore the hour of the day is 8 h. 2 m. 40 s. A.M.

7. Denote the arcs SP , SZ , and PZ (see Fig., p. 135) by z , p , and s .

Then in the triangle SPZ ,

$$z = 90^{\circ} - 15^{\circ} = 75^{\circ}, p = 90^{\circ}, s = 90^{\circ} - 42^{\circ} 21' = 47^{\circ} 39'.$$

Therefore in the polar triangle of SPZ , we have

$$\begin{aligned}Z &= 180^{\circ} - 75^{\circ} = 105^{\circ}, P' = 180^{\circ} - 90^{\circ} = 90^{\circ}, \\ S' &= 180^{\circ} - 47^{\circ} 39' = 132^{\circ} 21'.$$

$$\text{By Art. 144,} \quad \begin{array}{c} + \quad - \quad - \\ \cos p' = \cot S' \cot Z'. \end{array}$$

$$\log \cot S' = 9.9598$$

$$\log \cot Z' = \underline{9.4281}$$

$$\log \cos p' = \underline{9.3879}$$

$$\therefore p' = 75^{\circ} 51.6'.$$

Then in the triangle SPZ , we have

$$P = 180^{\circ} - p' = 104^{\circ} 8.4'.$$

To $104^{\circ} 8.4'$ corresponds 6 h. 56 m. 33.6 s. of time.

Therefore the hour of the day is 5 h. 3 m. 26.4 s. A.M.

USE OF THE TABLES.

Page 2.

- | | |
|--|---|
| 3. $\log 80 = 1.9031.$ | 12. Mantissa of 924 = .9657 |
| 4. $\log 6.3 = 0.7993.$ | $.61 \times 4 = \underline{2}$ |
| 5. $\log 298 = 2.4742.$ | $\therefore \log 92461 = 4.9659$ |
| 6. $\log .902 = 9.9552 - 10.$ | 13. Mantissa of 403 = .6053 |
| 7. Mantissa of 772 = .8876 | $.22 \times 11 = \underline{2}$ |
| $.3 \times 6 = \underline{2}$ | $\therefore \log .40322 = 9.6055 - 10$ |
| $\therefore \log .7723 = 9.8878 - 10$ | 14. Mantissa of 717 = .8555 |
| 8. Mantissa of 105 = .0212 | $.8 \times 6 = \underline{5}$ |
| $.6 \times 41 = \underline{25}$ | $\therefore \log .007178 = 7.8560 - 10$ |
| $\therefore \log 1056 = 3.0237$ | 15. Mantissa of 518 = .7143 |
| 9. Mantissa of 329 = .5172 | $.09 \times 9 = \underline{1}$ |
| $.4 \times 13 = \underline{5}$ | $\therefore \log 5.1809 = 0.7144$ |
| $\therefore \log 3.294 = 0.5177$ | 16. Mantissa of 103 = .0128 |
| 10. Mantissa of 520 = .7160 | $.65 \times 42 = \underline{27}$ |
| $.5 \times 8 = \underline{4}$ | $\therefore \log 1036.5 = 3.0155$ |
| $\therefore \log .05205 = 8.7164 - 10$ | 17. Mantissa of 866 = .9375 |
| 11. Mantissa of 200 = .3010 | $.76 \times 5 = \underline{4}$ |
| $.8 \times 22 = \underline{18}$ | $\therefore \log .086676 = 8.9379 - 10$ |
| $\therefore \log 20.08 = 1.3028$ | 18. Mantissa of 115 = .0607 |
| | $.07 \times 38 = \underline{3}$ |
| | $\therefore \log .11507 = 9.0610 - 10$ |

Page 4.

- | | |
|--|---|
| 4. Number corresponding to
1.8055 = 63.9. | 5. Number corresponding to
9.4487 - 10 = .281. |
|--|---|

$$\begin{array}{r}
 6. \quad 0.2165 \\
 \underline{.2148} = \text{mantissa of } 164 \\
 \frac{17}{27} = \quad \quad \quad 6
 \end{array}$$

\therefore Number corresponding = 1.646

$$\begin{array}{r}
 7. \quad 3.9487 \\
 \underline{.9484} = \text{mantissa of } 888 \\
 \frac{3}{5} = \quad \quad \quad 6
 \end{array}$$

\therefore Number corresponding = 8886

$$\begin{array}{r}
 8. \quad \text{Number corresponding to} \\
 \quad 2.7364 = 545.
 \end{array}$$

$$\begin{array}{r}
 9. \quad 8.1648 - 10 \\
 \underline{.1644} = \text{mantissa of } 146 \\
 \frac{4}{29} = \quad \quad \quad 1
 \end{array}$$

\therefore Number corresponding = .01461

$$\begin{array}{r}
 10. \quad 7.5209 - 10 \\
 \underline{.5198} = \text{mantissa of } 331 \\
 \frac{11}{13} = \quad \quad \quad 8
 \end{array}$$

\therefore Number corresponding = .003318

$$\begin{array}{r}
 11. \quad 4.0095 \\
 \underline{.0086} = \text{mantissa of } 102 \\
 \frac{9}{42} = \quad \quad \quad 21
 \end{array}$$

\therefore Number corresponding = 10221

$$\begin{array}{r}
 12. \quad 0.9774 \\
 \underline{.9773} = \text{mantissa of } 949 \\
 \frac{1}{4} = \quad \quad \quad 2
 \end{array}$$

\therefore Number corresponding = 9.492

$$\begin{array}{r}
 13. \quad 9.3178 - 10 \\
 \underline{.3160} = \text{mantissa of } 207 \\
 \frac{18}{21} = \quad \quad \quad 9
 \end{array}$$

\therefore Number corresponding = .2079

$$\begin{array}{r}
 14. \quad 1.6482 \\
 \underline{.6474} = \text{mantissa of } 444 \\
 \frac{8}{10} = \quad \quad \quad 8
 \end{array}$$

\therefore Number corresponding = 44.48

$$\begin{array}{r}
 15. \quad 7.0450 - 10 \\
 \underline{.0414} = \text{mantissa of } 110 \\
 \frac{36}{39} = \quad \quad \quad 9
 \end{array}$$

\therefore Number corresponding = .001109

$$\begin{array}{r}
 16. \quad 4.8016 \\
 \underline{.8014} = \text{mantissa of } 633 \\
 \frac{2}{7} = \quad \quad \quad 29
 \end{array}$$

\therefore Number corresponding = 63329

$$\begin{array}{r}
 17. \quad 8.1144 - 10 \\
 \underline{.1139} = \text{mantissa of } 130 \\
 \frac{5}{34} = \quad \quad \quad 1
 \end{array}$$

\therefore Number corresponding = 0.1301

$$\begin{array}{r}
 18. \quad 2.7015 \\
 \underline{.7007} = \text{mantissa of } 502 \\
 \frac{8}{9} = \quad \quad \quad 9
 \end{array}$$

\therefore Number corresponding = 502.9

Page 6.

3. $\log \tan 35^\circ 10' = 9.8479$

$$2.7 \times 9 = \underline{24}$$

$\therefore \log \tan 35^\circ 19' = 9.8503$

4. $\log \sin 61^\circ 50' = 9.9453$

$$.6 \times 8 = \underline{5}$$

$\therefore \log \sin 61^\circ 58' = 9.9458$

5. $\log \cot 12^\circ 30' = 0.6542$

$$5.9 \times 4 = \underline{24}$$

$\therefore \log \cot 12^\circ 34' = 0.6518$

6. $\log \cos 26^\circ 50' = 9.9505$

$$.6 \times 6 = \underline{4}$$

$\therefore \log \cos 26^\circ 56' = 9.9501$

7. $\log \tan 82^\circ 0' = 0.8522$

$$9.3 \times 3\frac{1}{3} = \underline{31}$$

$\therefore \log \tan 82^\circ 3' 20'' = 0.8553$

8. $\log \sin 55^\circ 10' = 9.9142$

$$.9 \times 1.8 = \underline{2}$$

$\therefore \log \sin 55^\circ 11.8' = 9.9144$

9. $\log \cos 30^\circ 40' = 9.9346$

$$.8 \times 2.5 = \underline{2}$$

$\therefore \log \cos 30^\circ 42.5' = 9.9344$

10. $\log \cot 48^\circ 0' = 9.9544$

$$2.5 \times 3\frac{3}{8} = \underline{9}$$

$\therefore \log \cot 48^\circ 3' 43'' = 9.9535$

Page 7.

3. 0.9164

$$0.9109 = \log \tan 83^\circ 0'$$

$$\frac{55}{10.5} = \underline{5.2}$$

$\therefore \text{Angle corresponding} = 83^\circ 5.2'$

4. 9.9238

$$9.9236 = \log \cos 33^\circ 0'$$

$$\frac{2}{.8} = \underline{2.5}$$

$\therefore \text{Angle corresponding} = 32^\circ 57.5'$

5. 9.8630

$$9.8629 = \log \sin 46^\circ 50'$$

$$\frac{1}{1.2} = \underline{0.8}$$

$\therefore \text{Angle corresponding} = 46^\circ 50.8'$

6. 0.2154

$$0.2127 = \log \cot 31^\circ 30'$$

$$\frac{27}{2.8} = \underline{9.6}$$

$\therefore \text{Angle corresponding} = 31^\circ 20.4'$

7. 9.2279

$$9.2251 = \log \cos 80^\circ 20'$$

$$\frac{28}{7.3} = \underline{3.8}$$

$\therefore \text{Angle corresponding} = 80^\circ 16.2'$

8. 9.4700

$$9.4669 = \log \cot 73^\circ 40'$$

$$\frac{31}{4.7} = \underline{6.6}$$

$\therefore \text{Angle corresponding} = 73^\circ 33.4'$

9. 9.1891

$$9.1863 = \log \sin 8^\circ 50'$$

$$\frac{28}{8} = \underline{3.5}$$

$\therefore \text{Angle corresponding} = 8^\circ 53.5'$

10. 0.0502

$$0.0481 = \log \tan 48^\circ 10'$$

$$\frac{21}{2.5} = \underline{8.4}$$

$\therefore \text{Angle corresponding} = 48^\circ 18.4'$

Page 8.

$$\begin{aligned}
 2. \log \sin 65^\circ 10' &= 9.9579 \\
 .5 \times 2 &= \underline{1} \\
 \log \sin 65^\circ 12' &= 9.9580 \\
 \therefore \log \csc 65^\circ 12' &= 0.0420
 \end{aligned}$$

$$\begin{aligned}
 3. \log \cos 80^\circ 0' &= 9.2397 \\
 7.3 \times 7.3 &= \underline{53} \\
 \log \cos 80^\circ 7.3' &= 9.2344 \\
 \therefore \log \sec 80^\circ 7.3' &= 0.7656
 \end{aligned}$$

$$\begin{aligned}
 4. \quad 9.5997 &= \log \cos. \\
 \underline{9.5978} &= \log \cos 66^\circ 40' \\
 \frac{19}{2.9} &= \underline{6.6} \\
 \therefore \text{Angle corresponding} &= 66^\circ 33.4'
 \end{aligned}$$

$$\begin{aligned}
 5. \quad 9.8112 &= \log \sin. \\
 \underline{9.8111} &= \log \sin 40^\circ 20' \\
 \frac{1}{1.4} &= \underline{0.7} \\
 \therefore \text{Angle corresponding} &= 40^\circ 20.7'
 \end{aligned}$$

Page 10.

$$\begin{aligned}
 3. \text{nat. sin } 3^\circ 30' &= .06105 \\
 .00029 \times 2 &= \underline{.00058} \\
 \text{nat. sin } 3^\circ 32' &= .06163 \\
 \text{mantissa of 616} &= .7896 \\
 7 \times .3 &= \underline{2} \\
 \therefore \log \sin 3^\circ 32' &= 8.7898
 \end{aligned}$$

$$\begin{aligned}
 4. \text{nat. cos } 88^\circ 10' &= .03199 \\
 .000291 \times 7 &= \underline{.00204} \\
 \text{nat. cos } 88^\circ 17' &= .02995 \\
 \text{mantissa of 299} &= .4757 \\
 14 \times .5 &= \underline{7} \\
 \therefore \log \cos 88^\circ 17' &= 8.4764
 \end{aligned}$$

$$\begin{aligned}
 5. \text{nat. tan } 2^\circ 20' &= .04075 \\
 .000291 \times 8.2 &= \underline{.00239} \\
 \text{nat. tan } 2^\circ 28.2' &= .04314 \\
 \text{mantissa of 431} &= .6345 \\
 10 \times .4 &= \underline{4} \\
 \therefore \log \tan 2^\circ 28.2' &= 8.6349
 \end{aligned}$$

$$\begin{aligned}
 6. \text{nat. tan } 50' &= .014545 \\
 .000291 \times 5\frac{14}{80} &= \underline{.001523} \\
 \text{nat. tan } 55' 14'' &= .016068
 \end{aligned}$$

$$\begin{aligned}
 \text{mantissa of 160} &= .2041 \\
 27 \times .68 &= \underline{18}
 \end{aligned}$$

$$\begin{aligned}
 \log \tan 55' 14'' &= 8.2059 \\
 \therefore \log \cot 55' 14'' &= 1.7941
 \end{aligned}$$

$$\begin{aligned}
 7. \quad 7.8702 \\
 \underline{.8698} &= \text{mantissa of } 741 \\
 \frac{4}{6} &= \underline{7}
 \end{aligned}$$

$$\begin{aligned}
 \text{Number corresponding} &= .007417 \\
 .007417 &= \text{nat. sin.} \\
 .005818 &= \text{nat. sin } 20' \\
 \underline{.001599} &= 5.497' = \underline{5' 29.8''} \\
 .0002909 &= \underline{5' 29.8''} \\
 \therefore \text{Angle corresponding} &= 25' 29.8''
 \end{aligned}$$

$$\begin{aligned}
 8. \quad 8.6150 \\
 \underline{.6149} &= \text{mantissa of } 412 \\
 \frac{1}{11} &= \underline{1} \\
 \text{Number corresponding} &= .04121
 \end{aligned}$$

$$\begin{aligned}
 .04121 &= \text{nat. cos.} \\
 .04071 &= \text{nat. cos } 87^\circ 40' \\
 \underline{.00050} &= 1.718' = \underline{1' 43.1''} \\
 .000291 &= \underline{1' 43.1''} \\
 \therefore \text{Angle corresp.} &= 87^\circ 38' 16.9''
 \end{aligned}$$

$$\begin{array}{r}
 9. \quad 8.2892 \\
 .2878 = \text{mantissa of} \quad 194 \\
 \hline
 \frac{14}{22} = \quad \quad \quad 64
 \end{array}$$

Number corresponding = .019464

$$.019464 = \text{nat. cot.}$$

$$.017455 = \text{nat. cot } 89^\circ 0'$$

$$\frac{.002009}{.0002905} = 6.916' = \quad 6' 55''$$

$$\therefore \text{Angle corresp.} = 89^\circ 53' 5''$$

$$\begin{array}{r}
 10. \quad 8.2131 = \text{cologarithm.} \\
 .2122 = \text{mantissa of} \quad 163 \\
 \hline
 \frac{9}{26} = \quad \quad \quad 35
 \end{array}$$

Number corresponding = .016335

$$.016335$$

$$.014545 = \text{nat. tan} \quad 50'$$

$$\frac{.001790}{.000291} = 6.151' = \quad 6' 9.1''$$

$$\quad \quad \quad 56' 9.1''$$

$$\therefore \text{Angle corresp.} = 89^\circ 3' 50.9''$$

Page 11.

$$3. \quad \text{nat. sin } 17^\circ 30' = .3007$$

$$.00028 \times 3 = \underline{.0008}$$

$$\therefore \text{nat. sin } 17^\circ 33' = .3015$$

$$4. \quad \text{nat. cos } 75^\circ 40' = .2476$$

$$.00029 \times 8.3 = \underline{.0024}$$

$$\therefore \text{nat. cos } 75^\circ 48.3' = .2452$$

$$5. \quad .7385$$

$$\frac{.7373}{.0012} = \text{nat. sin} \quad 47^\circ 30'$$

$$\frac{.0019}{.0019} = \quad \quad \quad 6.3$$

$$\therefore \text{Angle corresponding} = 47^\circ 36.3'$$

$$6. \quad .9280$$

$$\frac{.9272}{.0008} = \text{nat. cos} \quad 22^\circ 0'$$

$$\frac{.0011}{.0011} = \quad \quad \quad 7.3$$

$$\therefore \text{Angle corresponding} = 21^\circ 52.7'$$



LIBRARY OF CONGRESS



0 003 560 568 A

